



CORRECTION

Correction: Efficacy of Intravenous Treprostinil in High-Risk Single Ventricle Patients Undergoing Glenn Procedure

Xiaofeng Wang^{1,#}, Xingwei Chen^{2,#}, Shilin Wang¹, Xia Li¹, Zhongyuan Lu¹, Wenlong Wang¹ and Xu Wang^{1,*}

¹Department of Pediatric Intensive Care Unit, National Center for Cardiovascular Disease, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100037, China

²Department of Pharmacy, National Center for Cardiovascular Disease, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100037, China

*Corresponding Author: Xu Wang. Email: fwpicu@163.com

#These authors contributed equally to this paper

Received: 07 July 2025; Accepted: 07 July 2025; Published: 11 July 2025

In the article “Efficacy of Intravenous Treprostinil in High-Risk Single Ventricle Patients Undergoing Glenn Procedure” by Xiaofeng Wang, Xingwei Chen, Shilin Wang, Xia Li, Zhongyuan Lu, Wenlong Wang, Xu Wang (*Congenital Heart Disease*, 2024, vol. 19, no. 5, pp. 489–498. DOI:10.32604/chd.2024.054441, URL: <https://www.techscience.com/chd/v19n5/59166>), an error was identified in the reporting of statistical values in Table 1 and Table 4. Some *p*-values were incorrectly shown as “1.0”. The correct values are “>0.999”. The authors apologize for this oversight and present the corrected tables below.

Table 1: Demographic and surgical characteristics (revised).

Variable	Overall (n = 28)	Group 1 (n = 14)	Group 2 (n = 14)	<i>p</i> -Value
Age (years)	5.3 (0.9, 8)	6.7 (2.2, 10.5)	4.0 (0.7, 8)	0.205
Weight (kg)	15.3 (7.9, 21.3)	18.1 (9.8, 36.6)	8.6 (7.6, 19.5)	0.073
Heterotaxy syndrome (case, %)	4 (14)	3 (21)	1 (7)	0.589
Preoperative mPAP (mmHg)	17 ± 3	17 ± 3	18 ± 4	0.813
Cardiopulmonary bypass (min)	159 (86, 189)	165 (75, 189)	142 (98, 203)	0.554
Aortic cross-clamp (min)	83 (66, 113)	101 (63, 123)	80 (71, 92)	0.503
Other surgeries (case, %)	21 (75)	10 (71)	11 (79)	>0.999
Intraoperative mPAP (mmHg)	17 ± 3	17 ± 3	18 ± 3	0.292

Note: mPAP: mean pulmonary arterial pressure.

Table 4: Postoperative recovery (revised).

	Overall (n = 28)	Group 1 (n = 14)	Group 2 (n = 14)	<i>p</i> -Value
Mechanical ventilation duration (h)	47 (23, 122)	59 (22, 139)	28 (23, 123)	0.713
ICU length of stay (days)	9 (4, 18)	11 (4, 21)	7 (4, 13)	0.381
Postoperative length of stay (days)	21 (15, 33)	29 (19, 47)	18 (11, 22)	0.021
Duration of chest tube placement (days)	7 (6, 15)	7 (6, 18)	7 (6, 12)	0.762
Chest tube drainage volume (mL/kg)	31 (12, 88)	32 (10, 71)	29 (11, 101)	0.662
Thromboembolic events (cases, %)	3 (11)	2 (14)	1 (7)	>0.999
Tachycardia (cases, %)	12 (43)	6 (43)	6 (43)	>0.999
Renal replacement therapy (cases, %)	2 (7)	1 (7)	1 (7)	>0.999
Tracheal re-intubations (cases, %)	4 (14)	3 (21)	1 (7)	0.589
Tracheostomy (cases, %)	0 (0)	0 (0)	0 (0)	NA
Mortality (cases, %)	3 (11)	1 (7)	2 (14)	>0.999
Side effects (cases, %)	0 (0)	0 (0)	0 (0)	NA



The authors apologize for any inconvenience caused and affirm that the scientific conclusions remain unaffected. This correction has been reviewed and approved by the Editors-in-Chief of *Congenital Heart Disease*. The original article has been updated accordingly.