



ARTICLE

Bioactive Potential of *Calophyllum inophyllum*: Phytochemical Profiles, Biological Activities, and *In Silico* Pharmacokinetic Predictions

Luksamee Vittaya^{1,*} , Chakhriya Chalad¹ , Sittichoke Janyong² and Nararak Leesakul³

¹Department of General Education, Faculty of Science and Fisheries Technology, Rajamangala University of Technology Srivijaya, Sikao, Trang, 92150, Thailand

²Department of Marine Sciences and Environment, Faculty of Science and Fisheries Technology, Rajamangala University of Technology Srivijaya, Sikao, Trang, 92150, Thailand

³Division of Physical Science and Center of Excellence for Innovation in Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla, 90110, Thailand

*Corresponding Author: Luksamee Vittaya. Email: nokluksamee@hotmail.com or luksamee.v@rmutsv.ac.th

Received: 20 October 2025; Accepted: 28 November 2025; Published: 29 December 2025

ABSTRACT: *Calophyllum inophyllum* is a tropical plant that could have useful medicinal properties for pharmaceutical and cosmetic applications. The present study extracted the flower, fruit, leaf, twig, and bark of the plant by maceration in different organic solvents. The correlation between bioactive compounds and their biological activities was investigated, with emphasis on their therapeutic relevance through *in silico* pharmacokinetic predictions using SwissADME. Qualitative and quantitative analyses were conducted to determine the total phenolic, flavonoid, and saponin contents of the extracts. Spectral analysis of the extracts revealed –OH, C=O, C=C, and C–H functional groups. The antioxidant activity of the extracts was evaluated by colorimetric DPPH and ABTS assays. Antibacterial activity was also determined, along with cytotoxicity and anticancer potential. The plant part and solvent used for extraction affected the active compounds collected. Methanolic bark extract presented the highest phenolic and flavonoid contents, whereas hexane bark extract presented the highest saponin content. These results correlated with the antioxidant and antibacterial activities of the extracts. The strong correlation between antioxidant activity and total phenolic contents indicated that phenolic compounds were the dominant contributors to antioxidant activity. Methanolic bark extract showed significant DPPH scavenging activity ($IC_{50} = 0.004$ mg/mL) while hexane bark extract produced the largest zone of inhibition against *Staphylococcus aureus* (17.96 ± 0.00 mm). The lowest minimum inhibitory concentration (<0.098 mg/mL) was shown against *Bacillus cereus* by 11 out of the 15 extracts. Selective cytotoxicity was observed against cancer cells, especially lung cancer cells. *In silico* ADME analysis of a previously reported flower extract showed favorable pharmacokinetic properties for compounds such as eugenol and caryophyllene oxide. High gastrointestinal absorption and blood-brain barrier permeability suggested good oral bioavailability and central nervous system potential. Most extracted compounds met Lipinski's Rule with minimal cytochrome P450 inhibition, indicating good drug-likeness. These findings highlight the promise of *C. inophyllum* as a source of natural antioxidants and antibacterial agents with therapeutic potential.

KEYWORDS: *Calophyllum inophyllum*; antibacterial; antioxidant; mangrove; metabolite; ADME

1 Introduction

The uncertain climatic conditions that are affecting human settlement and food supplies are also contributing to the emergence of new diseases [1]. Currently, synthetic antibiotics are widely used to manage diseases [2], but the adoption of natural products into disease management might be a more promising approach to the prevention of emerging diseases. Medicinal plants and herbs have attracted



significant attention as potential alternatives to synthetic antibiotics, and recent research has focused on identifying plants that thrive in fluctuating environmental conditions, such as those found in mangrove or coastal forests. These plants not only survive but thrive, adapting to harsh conditions by producing bioactive compounds to protect themselves from harmful environmental factors. Compounds from the mangrove plant *Acanthus ebracteatus* were shown to exhibit antioxidant and biological activities [3], compounds from *Pluchea indica* have shown pharmacological properties [4], and *Calophyllum inophyllum* has been widely studied for its potential to contribute to pharmaceutical and cosmetic applications [5]. In addition, mangrove plants such as *Rhizophora mucronata* and *Rhizophora apiculata* are cooked and eaten for their medicinal properties [6,7]. Therefore, the discovery of bioactive substances from these plants supports local traditional wisdom and extends the application and utilization of plants and herbs in medicine.

C. inophyllum is a mangrove plant and consequently endures constantly changing environmental conditions. Researchers have extracted various parts of this plant, isolating and investigating several significant bioactive compounds from the seed, fruit, leaf, and bark [5,8,9]. However, the isolation and purification processes are time-consuming and require substantial quantities of solvent and advanced analytical tools. Preliminary phytochemical screening provides a more efficient and rapid method of classifying bioactive substances, enabling quantitative studies to identify potential applications. Understanding the biological activities of different parts of *C. inophyllum* can assist in the identification of phytochemicals that can provide pharmaceutical or medical alternatives to synthetic chemicals and antibiotics. Moreover, this approach would accelerate the use of extracts by shortening the lengthy process of isolating specific compounds.

Recently, our research group conducted a rare study of the flower of *C. inophyllum*. The results showed that macerated extracts of *C. inophyllum* flower [10] yielded a broader range of active ingredients than distilled extracts [11,12]. Gas chromatography-mass spectrometry revealed various important bioactive compounds in the flower extracts. The chemical composition of the extracts was found to depend on the solvent used for extraction [10]. The polarity of the solvent and solute interactions may explain the greater antibacterial activity of the hexane extract of *C. inophyllum* flower compared to the ethyl acetate and methanol extracts. The bioactive compounds detected in these extracts included α -copaene, α -muurolene, β -muurolene, γ -cadinene, δ -cadinene, caryophyllene oxide, eugenol, palmitic acid, and phytol, each exhibiting different biological activities. To further explore the pharmaceutical potential of these compounds, an *in silico* evaluation of their molecular properties was undertaken in the present study. Absorption, distribution, metabolism, and excretion (ADME) were studied to predict the physicochemical properties of the compounds, their lipophilicity, water solubility, drug-likeness, pharmacokinetics, and medicinal chemistry suitability. This analysis was expected to facilitate the assessment of their potential as therapeutic agents [13]. The ADME computational approach is particularly valuable for determining the oral bioavailability, solubility, and potential drug interactions of bioactive compounds, thereby streamlining the identification of promising drug candidates from natural sources.

This research aims to determine the contents of phenolics, flavonoids, and saponins in *C. inophyllum* fruit, leaf, twig, and bark extracts for comparison with the contents of the previously investigated flower extracts. The ethnomedicinal properties of the plant were examined using both qualitative and quantitative analyses, and the plant parts were screened for pharmaceutically important bioactive substances. Each plant part was extracted using three solvents of varying polarity: hexane, ethyl acetate, and methanol. Fourier transform infrared (FTIR) and ultraviolet-visible (UV-Vis) spectroscopy were employed to identify the functional groups and chromophores present in the bioactive compounds. To the best of our knowledge, based on a thorough literature review, this correlation-based study is the first to quantify and compare the phytochemicals and the antioxidant and antibacterial activities of extracts obtained from various parts

of *C. inophyllum* with different solvents. The cytotoxicity and anticancer potential of the extracts were also evaluated. This study combines solvent–part interaction analysis with *in silico* ADME prediction to provide a clearer understanding of the influence of extraction conditions on the bioactive composition and pharmacological potential of *C. inophyllum*. The findings will be valuable for practical applications and for identifying the key factors influencing biological activity across different plant parts. Overall, the results highlight the potential of *C. inophyllum* as a natural source of bioactive ingredients for food, cosmetic, and medicinal applications.

2 Material and Methods

2.1 Chemicals and Reagents

The extracting solvents used in this study (hexane, ethyl acetate, and methanol) were of analytical grade, obtained from RCI Labscan (Bangkok, Thailand) and used without further purification. Dimethyl sulfoxide (DMSO) was from Merck (Darmstadt, Germany). Folin-Ciocalteu reagent and aluminum chloride were from Loba Chemie (Mumbai, India). Gallic acid, rutin, escin, DPPH, ABTS⁺, and p-iodonitrotetrazolium chloride were purchased from Sigma Aldrich (MO, USA). Mueller Hinton broth (MHB) was from Difco, Becton Dickinson (MD, USA). Petroleum ether was supplied by Fisher Scientific (Leicester, UK), and vanillin solution was obtained from Himedia (Mumbai, India).

2.2 Plant Materials

Plants were collected from Trang Province, Thailand, at the coordinates 7°52'20'' N, 99°30'72'' E. A voucher specimen (BKF 194811) was deposited at the Forest Herbarium, Bangkok, Department of National Parks, Wildlife and Plant Conservation, Thailand (Fig. 1). Fresh flowers, fruits, leaves, and twigs were collected in September, and bark samples were collected in October from mature plants. All plant materials were properly cleaned to remove debris and shade-dried at room temperature for further experiments.



Figure 1: The flower, fruit, leaf, twig, and bark of *Calophyllum inophyllum*.

2.3 Extraction Procedure

Each dried plant part (1000 g) was extracted by maceration using hexane, ethyl acetate, and methanol sequentially at a ratio of 1:7 (plant dry weight–solvent volume) for one week. The obtained extracts were evaporated to dryness, using a rotary evaporator at 45°C. The concentrated extracts were stored in amber glass bottles at 4°C in a refrigerator until further analysis. The yields of flower, fruit, leaf, twig, and bark extracts were reported. All extracts were refrigerated prior to the quantitative analysis of phytochemical contents and antioxidant and antibacterial activities.

2.4 Phytochemical Analysis

2.4.1 Qualitative Screening

The preliminary phytochemical screening of *C. inophyllum* extracts was carried out according to the method described by Vittaya et al. [10]. To test for anthraquinone, the crude extract was shaken with a 10% solution of sulfuric acid (H_2SO_4) and heated for 5 min. The filtered extract was then extracted three times with 5 mL of chloroform. The extract was then introduced into a 3% ammonia solution (NH_3 dilution). The appearance of a pink color in the ammonia layer indicated the presence of anthraquinone. To test for terpenoids, the crude extract was dissolved a few times in 3 mL of petroleum ether and collected by filtration. Slowly, 2 mL of chloroform and concentrated sulfuric acid (H_2SO_4) were added to the filtrate. A brown-ring between two layers indicated the presence of terpenoids. To test for flavonoids, the crude extract was mixed with 95% ethanol, and 2–3 small pieces of magnesium were introduced. The mixture was then filtered, and a few drops of concentrated hydrochloric acid (HCl) were added. The formation of a yellowish-orange precipitate indicated the presence of flavonoids. To test for saponins, 5 mL of water was added to the crude extract. The mixture was warmed, and filtered. Subsequently, 2–3 mL of water was added to the filtrate that was then shaken vigorously for 10 min. The appearance of foam indicated the presence of saponins. To test for phenolics, a few drops of 1% ferric chloride (FeCl_3) solution were added to the crude extract, which was shaken with water. A dark brown or blackish-green color indicated the presence of phenolics. To test for alkaloids, the crude extract was shaken in 15 mL of 2% sulfuric acid (H_2SO_4) and warmed. Dragendorff's reagent was then added to the mixture. The formation of an orange-brown precipitate indicated the presence of alkaloids.

2.4.2 Quantitative Analysis

Determination of Phenolic Content. The total phenolic content (TPC) of the extracts was determined spectroscopically using the Folin-Ciocalteu colorimetric method [14] with gallic acid as a standard. The absorbance of the reaction mixture was recorded at 765 nm using a UV-visible spectrophotometer (U-1800 spectrophotometer, Hitachi High-Tech Science Corp., Tokyo, Japan). The TPC for each sample was calculated using the standard curve of gallic acid at concentrations from 100–600 $\mu\text{g/mL}$, and expressed in terms of gallic acid equivalents (mg gallic acid/g crude extract). The data were calculated according to the equation $y = 0.0040x - 0.0086$, $R^2 = 0.9977$.

Determination of Flavonoid Content. The total flavonoid content (TFC) of the extracts was determined using the aluminium chloride method [14] with rutin as a standard. The absorbance was recorded at 510 nm using a UV-visible spectrophotometer, with measurements taken against a blank sample. TFC was calculated from a standard calibration curve prepared with rutin solutions (100–600 $\mu\text{g/mL}$) and expressed in terms of rutin equivalents (mg rutin/g crude extract). The data were calculated according to the equation $y = 0.0016x + 0.0002$, $R^2 = 0.9992$.

Determination of Saponin Content. The total saponin content (TSC) of the extracts was determined according to the method described by Vittaya et al. [10] and Senguttuvan et al. [15], with a modification involving a reduced extract volume. In this method, 0.2 mL of the extract was mixed with 0.5 mL of a 0.8% (w/v) vanillin solution, followed by the addition of 5 mL of 72% (v/v) sulphuric acid. The mixture was allowed to stand for 1 min, then incubated at 70°C for 10 min. After incubation, the solution was rapidly cooled in an ice-water bath to room temperature. The absorbance was measured at 560 nm using a UV-visible spectrophotometer. Methanol was used as the control, while escin served as the reference standard for comparison. The TSC was expressed in terms of escin equivalent (mg escin/g crude extract) through the calculation curve of escin ($y = 0.0007x + 0.0254$), which was calibrated alongside the standard curve of escin at concentrations of 100–1000 µg/mL. The experiment was performed in triplicate, and the data were expressed as means \pm standard deviation with a linearity R^2 equal to 0.9918.

2.5 Structural Characterization

2.5.1 UV-Visible Absorption Spectroscopy

UV-Visible spectral analysis by Hitachi UV-Visible spectrophotometer (U-5100) with UV solutions 4.2 program (Tokyo, Japan) was done in the range of 200–800 nm with a resolution of 1 nm. One milliliter of sample was pipetted into a test tube and analyzed at room temperature. The solvent used to extract each plant part was utilized as a blank.

2.5.2 Fourier Transform Infrared Spectroscopy (FTIR) Analysis

FTIR spectra were recorded with the PerkinElmer BX FTIR spectrophotometer (CT, USA). All samples were prepared by mixing approximately 1 mg of *C. inophyllum* extract with 3 mg of dry potassium bromide (KBr), using a mortar and pestle. The resulting homogeneous mixture was then pressed into a disc for analysis. The analysis covered a wavelength range of 4000–400 cm^{-1} at a temperature of 25°C. The spectra were recorded in transmittance mode, with the absorbance of each sample expressed as a percentage of transmittance (%T).

2.6 Antioxidant Activity Assays

2.6.1 DPPH Radical Scavenging

DPPH radical scavenging activity was determined using spectroscopy, based on the modified method described by Vittaya et al. [6]. Briefly, a stock solution of extract or standard was prepared at a concentration of 1 mg/mL, and then 0.5 mL of this solution was mixed with 0.5 mL of 0.15 M DPPH solution. The reaction mixture was left to incubate in the dark at room temperature for 30 min. The absorbance of the reaction was measured at 517 nm against the blank. The percentage of free radical scavenging activity of sample was calculated using Eq. (1):

$$\% \text{ Free radical scavenging activity} = [1 - (A_{\text{sample}} - A_{\text{sample blank}})/A_{\text{control}}] \times 100 \quad (1)$$

where A_{sample} is the absorbance of the test sample with DPPH solution, $A_{\text{sample blank}}$ is the absorbance of the test sample only and A_{control} is the absorbance of the DPPH solution.

2.6.2 ABTS Radical Scavenging

ABTS radical scavenging activity was determined following the modified method described by Vittaya et al. [6]. In brief, the ABTS⁺ working solution was diluted to achieve an absorbance of 0.700 ± 0.025 unit at 734 nm. In a centrifuge tube, 0.9 mL of diluted ABTS⁺ working solution was added to 0.1 mL of standard

or sample extract. The reaction mixture was incubated at room temperature for 6 min. After incubation, absorbance was recorded at 734 nm. The percentage of free radical scavenging activity for each sample was determined using Eq. (2):

$$\% \text{ Free radical scavenging activity} = [(A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}] \times 100 \quad (2)$$

where A_{control} is the absorbance of the extract without ABTS⁺ solution and A_{sample} is the absorbance of the extract with ABTS⁺ solution.

2.7 Biological Activity Analysis

2.7.1 Antibacterial Screening

The antibacterial activity of all extracts was preliminarily assessed in triplicate using the paper disc diffusion method [16]. Seven bacterial species were tested: *Bacillus cereus* TISTR 687, *Staphylococcus aureus* TISTR 1466, *Staphylococcus epidermidis* TISTR 518, *Escherichia coli* TISTR 780, *Salmonella typhi* TISTR 292, *Klebsiella pneumoniae* TISTR 1843, and *Pseudomonas aeruginosa* TISTR 781. These strains were obtained from the National Center for Genetic Engineering and Biotechnology (BIOTEC), Pathumthani, Thailand. For the disc diffusion assay, the bacterial cultures were incubated in tryptic soy broth at 35°C for 24 h, and their turbidity was measured against 0.5 McFarland standards (1×10^8 colony forming units/mL). With a sterile cotton swab, the bacteria were swabbed over the surface of the media (tryptic soy agar) and allowed to solidify. After incubation at 35°C for 24 h, the inhibition zones were measured in millimetres. To provide negative controls, the fifteen extracts were dissolved in dimethyl sulfoxide (DMSO) to prepare stock solutions of 100 mg/mL. Ten microliters of the solutions were placed on sterile filter paper discs (6 mm in diameter, Whatman No. 1 filter paper). Gentamicin was the positive control.

2.7.2 Determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

Stock solutions of the seven model pathogenic bacteria were prepared, and all tested inoculums were produced as 1×10^6 CFU/mL in Mueller Hinton broth (MHB) using a modified method [17]. All extracts and control stock solutions were prepared at an initial concentration of 100 mg/mL and serially diluted two-fold with MHB to obtain a concentration range from 0.01–50,000 µg/mL. In 96-well plates, 50 µL of each diluted concentration was mixed with 50 µL of MHB. The standardized inoculums at 1×10^6 CFU/mL were then introduced into each well and gently mixed by multichannel auto-pipette to produce a final concentration of 5×10^5 CFU/mL. The plates were then covered with sterile plate sealers. After incubation at 35°C for 24 h, the turbidity of the solutions was measured. Bacterial growth was confirmed by the color change from yellow to pink of p-iodonitrotetrazolium chloride (INT) dye added to the wells. The MIC was defined as the lowest extract concentration that completely inhibited bacterial growth by preventing the color change. The quadruplicate tests included a negative growth control of DMSO (CLSI, 2019). After the broth microdilution tests, the MBC was determined by the drop plate technique. Briefly, a volume of 10 µL was removed from the microtiter plate wells where no growth was observed, and then inoculated onto the surface of HMA plates. The plates were incubated at 35°C for 24 h, and the MBC was taken to be the lowest concentration of the substance at which no colonies formed under these conditions.

2.7.3 Cytotoxicity and Anticancer Assays

The anticancer activity of *C. inophyllum* extracts was assessed using the resazurin microplate assay (REMA) against three cancer cell lines: oral cavity cancer (KB), breast cancer (MCF7), and small cell lung

cancer (NCI-H187). The cytotoxicity of the extracts was also evaluated against African green monkey kidney (Vero) cells as a normal cell line control. The positive controls used for comparison were Ellipticine, Doxorubicin, and Tamoxifen in dimethyl sulfoxide (DMSO). For the anticancer screening, the extracts were tested at various concentrations to determine the percentage of cytotoxicity and the IC₅₀ values. The screening was performed at the National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), Pathumthani, Thailand. The threshold for cytotoxicity was set at 50%, with values below 50% indicating non-cytotoxicity and values above 50% indicating cytotoxicity. IC₅₀ values were calculated for extracts that demonstrated cytotoxicity against the cancer cell lines [18].

2.8 In Silico ADME Profile

The primary physicochemical, lipophilicity, water solubility, pharmacokinetic, drug-likeness, and medicinal chemistry properties of phytochemical compounds identified from the extracts of *C. inophyllum* flower were evaluated for ADME (absorption, distribution, metabolism, and excretion) prediction. The initial step involved accessing the PubChem database (<https://pubchem.ncbi.nlm.nih.gov/>) to obtain canonical SMILE (Simplified Molecular Input Line Entry) system representations of each compound. Subsequently, these SMILES were used for ADME prediction using the SwissADME online server (<http://www.swissadme.ch/>) [13].

2.9 Statistical Analysis

The data were analysed using IBM SPSS software version 25.0 (IBM Corp., Armonk, NY, USA) and expressed as means \pm standard deviation (SD) of triplicate determinations ($n = 3$). The $p < 0.05$ was considered to be statistically significant. Statistical analysis was carried out using a one-way analysis of variance (ANOVA), followed by Duncan's new multiple range test. Two-way ANOVA was used to investigate the interaction between plant parts (flower, fruit, leaf, twig, and bark) and the three organic solvents (hexane, ethyl acetate, and methanol). Pearson correlation was performed to evaluate the correlation between phytochemical compositions (TPC, TFC, and TSC) and antioxidant activity, as well as antibacterial activities against the seven model pathogenic bacteria. Principal component analysis (PCA) was conducted using a correlation matrix to evaluate the contribution of variables (assay) and factors (plant part and solvent) to the characteristics of *C. inophyllum* extracts.

3 Results and Discussion

3.1 Plant Extraction and Phytochemical Screening

Plants produce primary and secondary metabolites. Secondary metabolites play an important role in the growth of plants that survive changing terrain and climate. They include phenolics, flavonoids, alkaloids, and terpenoids. These compounds are produced as part of defense mechanisms and exhibit complex structures which depend on diverse polarities and vary across species [19]. The polarity of organic solvents used to extract these bioactive compounds has a significant impact on extraction efficiency. In this study, *C. inophyllum* flower, leaf, fruit, twig, and bark were extracted by maceration in hexane, ethyl acetate, and methanol. Fifteen extracts were obtained from the different plant parts. The data were presented in Table 1, and these extracts were investigated in phytochemical and biological analyses.

Methanol extracted the highest yields from all plant parts, followed by ethyl acetate then hexane, except in the case of fruit. This result indicated that most of the compounds extracted from the various plant parts were highly polar, as methanol is a highly polar solvent and, according to the principle of 'like dissolves like,' polar solvents tend to dissolve polar compounds. The solubility of organic compounds

is influenced by factors such as the number of hydroxyl groups present, molecular weight, and carbon chain length [20,21]. Methanol contains hydrogen bonds, which increase the solubility of both primary and secondary metabolites. The type of phytochemicals extracted also depends on the plant material. The compounds present in plant metabolites can vary from simple to highly polymerized substances such as phenolic acid, flavonoids, alkaloids, saponins, terpenoids, and anthraquinones [6,22]. Primary metabolites, such as carbohydrates and proteins, and secondary metabolites, such as phenolic compounds and flavonoids, are more soluble in methanol. While ethyl acetate is a medium-polar solvent, it can extract compounds in the same polar group but less effectively than methanol. In contrast, hexane is a non-polar solvent, and can extract compounds such as hydrocarbons, lipids, steroids, and fats [20,22]. The different yields obtained with these organic solvents (Table 1) reflect their different polarities.

Table 1: Yields of various extracts of *C. inophyllum*.

Plant	Parts	Solvents	Yield (g)	%
<i>C. inophyllum</i>	Flower*	Hexane	6.73	1.64
		Ethyl acetate	23.30	5.68
		MeOH	241.46	58.89
	Fruit	Hexane	221.64	11.31
		Ethyl acetate	126.02	6.43
		MeOH	378.37	19.30
	Leaf	Hexane	47.33	2.37
		Ethyl acetate	96.59	4.83
		MeOH	234.56	11.73
	Twig	Hexane	42.77	2.04
		Ethyl acetate	68.73	3.27
		MeOH	164.53	7.83
	Bark	Hexane	87.62	6.44
		Ethyl acetate	138.17	10.16
		MeOH	320.55	23.57

*Data from the flower extracts are reused in this article by permission from Elsevier (License No. 5795780615723) [10].

The results presented in Table 2 revealed that hexane extracts of all the plant parts contained only terpenoids. In contrast, ethyl acetate extracts exhibited the presence of terpenoids, flavonoids, and phenolics, while methanol extracts showed the most diverse range of metabolites, including terpenoids, flavonoids, saponins, phenolics, alkaloids, and anthraquinones. All the extracted phytochemicals were found to some degree among the leaf, twig, and bark extracts. Anthraquinones are structurally aromatic organic compounds which consist of three benzene rings bonded to each other, and they are found in traditional medicines mainly used as laxatives [23]. Terpenoids are small secondary metabolites found in many plants, and are known for their diverse biological activities that include antioxidant and antimicrobial properties [24]. Flavonoids, a chemically distinct group of polyphenols with conjugated aromatic systems, exhibit antioxidant, antibacterial, anti-inflammatory, and antiallergic activities [25]. Certain flavonoids, including quercetin and rutin, play medicinal roles in human health, showing anti-inflammatory, antiviral, and anticancer activities [26,27]. Saponins are glycosides that have amphiphile properties and are soluble in both water and lipids. Previous studies have identified saponins in *Moringa oleifera* leaves, where they demonstrated cholesterol-lowering properties [28]. In addition, saponins have been shown to inhibit the growth of human gastric cancer cells [29]. Phenolics, a group of phytochemicals with one or more aromatic rings and hydroxyl groups, include phenolic acids, flavonoids, tannins, and coumarins, and are recognized for their antioxidant, antimicrobial, and anticancer activities [30]. Finally, alkaloids are organic compounds that contain nitrogen within the molecule. The functions of alkaloids in plants are assumed to be as a

source of nitrogen for protein production, control of growth or germination of seeds, and protection from insects [31].

Table 2: Qualitative phytochemical analysis of various extracts of *C. inophyllum*.

Plant Constituent	Verification Method	Observations	<i>C. inophyllum</i>														
			Flower			Fruit			Leaf			Twig			Bark		
			H	E	M	H	E	M	H	E	M	H	E	M	H	E	M
Anthraquinone	Borntrager’s test	Formation of a rose-pink color	—		+	—	—	+	—	—	+	—	—	+	—	—	+
Terpenoid	Salkowski’s test	Reddish brown ring	+	+	—	+	+	+	+	—	—	+	—	—	+	—	—
Flavonoid	Reduction of metal	Formation of a cherry color	—	+	+	—	—	—	—	—	+	—	—	+	—	+	+
Saponin	Forth test	Formation of a stable form	—	—	+	—	—	+	—	—	+	—	—	+	—	—	+
Phenolic	Ferric chloride test	Green-bluish, dark green	—	+	+	—	+	+	—	+	+	—	—	+	—	—	+
Alkaloid	Dragendorff’s test	Formation of an orange-yellow precipitate	—	—	—	—	—	—	—	—	+	—	—	+	—	—	+

+: present and —: absent. H: hexane, E: ethyl acetate, and M: methanol.

3.2 Structural Characterization

3.2.1 UV-Visible Absorption Spectroscopy

UV-vis spectra of flower, leaf, fruit, twig, and bark extracts of *C. inophyllum* exhibited absorbance bands in the 200–400 nm range (Fig. 2). The leaf extracts also produced an absorption band in the 600–700 nm range, which corresponds to the absorbance of chlorophyll [32]. The first photoabsorption band of phenolic compounds, around 265–315 nm, is typically attributed to their aromatic structure. Kumar and Goel [30] reported that phenolic compounds absorb in the UV region of 225–330 nm. The solvent used for extraction had a greater effect on absorption than the plant part extracted. The absorption peaks shifted due to the varying polarity of the solvents, which affects the solubility and extraction efficiency of phenolic compounds. The twig and bark extracts showed stronger absorbance in the UV region compared to the leaf extracts. This suggests that the twig and bark parts of *C. inophyllum* contained a higher concentration of bioactive compounds, especially phenolics, which contribute significantly to UV absorption. These results are consistent with the observed total phenolic contents (TPC) (Table 3). The lower TPC observed in hexane extracts can be attributed to the non-polar nature of hexane. However, ethyl acetate and methanol extracts demonstrated higher TPC, indicating a greater presence of hydroxyl (–OH) groups, which are known for their strong hydrogen bonding and intermolecular interactions, which enhance the stability and activity of various phytochemicals [30]. The UV absorption spectra for the flower and fruit extracts of *C. inophyllum* displayed solvent-dependent variations, with methanol and ethyl acetate extracts showing stronger absorption than hexane extracts. These results indicated that the polar solvents extracted these bioactive compounds more efficiently from *C. inophyllum*.




Figure 2: Absorption spectra of *C. inophyllum* extracts from different plant parts: flower (A), fruit (B), leaf (C), twig (D), and bark (E), extracted with hexane, ethyl acetate (EtOAc), and methanol (MeOH).

Table 3: Total phenolic content (TPC), total flavonoid content (TFC), total saponin content (TSC), and radical scavenging activity of *C. inophyllum* extracts. Two-way analysis of variance showed the effects of plant parts (PP), solvent (S), and their interaction.

Parts	Solvents	TPC (mg GAE/g CE)	TFC (mg RU/g CE)	TSC (mg EE/g CE)	IC ₅₀ DPPH (mg/L)	IC ₅₀ ABTS (mg/L)					
Flower*	Hexane	0.19 ± 0.02 ^{ij}	0.91 ± 0.18 ^{gh}	5.05 ± 0.79 ^{bc}	0.343 ± 0.003 ⁱ	0.117 ± 0.001 ^g					
	Ethyl acetate	0.84 ± 0.07 ^c	1.48 ± 0.11 ^{ef}	3.14 ± 0.55 ^{d-f}	0.014 ± 0.000 ^{a-c}	0.077 ± 0.004 ^{ef}					
	Methanol	0.57 ± 0.05 ^{c-g}	1.03 ± 0.07 ^g	2.99 ± 0.41 ^{ef}	0.030 ± 0.001 ^{b-d}	0.047 ± 0.003 ^c					
Fruit	Hexane	0.18 ± 0.02 ^{ij}	1.34 ± 0.07 ^f	1.31 ± 0.35 ^h	0.502 ± 0.002 ^j	0.211 ± 0.010 ⁱ					
	Ethyl acetate	0.72 ± 0.06 ^{c-e}	1.30 ± 0.13 ^f	1.76 ± 0.45 ^{gh}	0.141 ± 0.002 ^g	0.071 ± 0.000 ^e					
	Methanol	0.20 ± 0.02 ^{ij}	0.70 ± 0.06 ^h	2.06 ± 0.51 ^{f-h}	0.109 ± 0.001 ^f	0.058 ± 0.002 ^d					
Leaf	Hexane	0.15 ± 0.02 ^j	1.96 ± 0.19 ^{bc}	5.35 ± 0.63 ^{bc}	0.852 ± 0.045 ^k	0.172 ± 0.003 ^h					
	Ethyl acetate	0.65 ± 0.28 ^{d-f}	2.03 ± 0.19 ^b	2.30 ± 0.32 ^{f-h}	0.042 ± 0.001 ^d	0.059 ± 0.004 ^d					
	Methanol	0.77 ± 0.06 ^{cd}	1.75 ± 0.14 ^{cd}	2.60 ± 0.37 ^{fg}	0.032 ± 0.000 ^{cd}	0.045 ± 0.003 ^c					
Twig	Hexane	0.33 ± 0.03 ^{hi}	0.89 ± 0.06 ^{gh}	5.19 ± 0.89 ^{bc}	0.188 ± 0.005 ^h	0.076 ± 0.003 ^{ef}					
	Ethyl acetate	0.69 ± 0.06 ^{c-e}	1.68 ± 0.17 ^{de}	3.80 ± 0.71 ^{de}	0.027 ± 0.000 ^{b-d}	0.045 ± 0.001 ^c					
	Methanol	1.48 ± 0.13 ^b	2.86 ± 0.12 ^a	4.24 ± 0.76 ^{cd}	0.027 ± 0.000 ^{b-d}	0.048 ± 0.005 ^c					
Bark	Hexane	0.42 ± 0.02 ^{gh}	0.24 ± 0.01 ⁱ	7.89 ± 1.01 ^a	0.132 ± 0.005 ^g	0.080 ± 0.000 ^f					
	Ethyl acetate	0.51 ± 0.03 ^{fg}	0.81 ± 0.06 ^{gh}	5.66 ± 0.89 ^b	0.062 ± 0.002 ^e	0.051 ± 0.002 ^c					
	Methanol	1.81 ± 0.14 ^a	3.07 ± 0.11 ^a	7.17 ± 0.72 ^a	0.004 ± 0.000 ^a	0.063 ± 0.001 ^d					
BHT					0.011 ± 0.000 ^{ab}	0.017 ± 0.001 ^b					
Vit C					0.004 ± 0.000 ^a	0.002 ± 0.000 ^a					
Two-Way ANOVA											
Variable	df	F	P	F	P	F	P	F	P	F	P
PP	4	51.830	<0.001	80.876	<0.001	73.781	<0.001	753.272	<0.001	342.580	<0.001
S	2	209.899	<0.001	160.703	<0.001	24.045	<0.001	4559.042	<0.001	2071.829	<0.001
PP × S	8	50.424	<0.001	126.575	<0.001	4.484	0.001	577.928	<0.001	241.216	<0.001
Error	30										

*Data from the flower extracts are reused in this article by permission from Elsevier (License No. 5795780615723) [10]. Data were presented as means ± SD ($n = 3$). Means ± SD in a column superscripted with different lowercase letters denote significant differences ($p < 0.05$).

3.2.2 FTIR Spectroscopy

The FTIR spectra of the extracts of *C. inophyllum* flower are presented with peak assignments (Fig. 3). The O–H stretching peaks at $3000\text{--}3500\text{ cm}^{-1}$, $2700\text{--}3000\text{ cm}^{-1}$, and $500\text{--}1000\text{ cm}^{-1}$ correspond to the stretching oscillations of alcohol or carboxylic acid groups which overlap the stretching vibration attributed to water. The elongation bands of C–H on the aromatic ring are present at $3500\text{--}4000\text{ cm}^{-1}$. Additionally, the band at $2700\text{--}3000\text{ cm}^{-1}$ corresponds to C–H stretching in aliphatic hydrocarbons, which indicates the presence of fatty acids, lipids, and possibly essential oils [33]. The sp^2 C–H stretching band in this range is associated with aromatic compounds, which may include flavonoids and phenolic acids. Moreover, the flexion band around $1500\text{--}1800\text{ cm}^{-1}$ corresponds to C=C bending in aromatic rings, C=C stretching in alkenes, or C=O stretching in carboxylic acid derivatives, suggesting the presence of carbonyl compounds such as terpenoids or aldehydes [34]. The oscillation of amine stretching and C–N bending vibrations were detected at $1000\text{--}1500\text{ cm}^{-1}$ [35]. The peak at $1000\text{--}1200\text{ cm}^{-1}$ is the elongation band of C–O stretching of methoxy ($-\text{OCH}_3$) in alkyl aryl ether or ester compounds. It is possible that the absorption peak at $500\text{--}580\text{ cm}^{-1}$ may correspond to the stretching of C–Br in alkyl halide or the presence of substitution groups on the benzene ring [36]. The observed peaks confirmed the presence of the functional groups –OH, C=O, C–H, C=C, C–O, and N–H, and support the findings of previous studies on mangrove plants and their phytochemicals [37]. These results also support the findings of previous reports which demonstrated that *C. inophyllum* extracts contained phenolic and flavonoid components with potential biological activities [5,10].




Figure 3: FTIR spectra of *C. inophyllum* extracts from various plant parts: flower (A), fruit (B), leaf (C), twig (D), and bark (E), extracted using hexane (H), ethyl acetate (E), and methanol (M).

3.3 Phytochemical Study of *C. inophyllum*

The TPC, TFC, and TSC of *C. inophyllum* extracts ranged from 0.19 to 1.81 mg GAE/g crude extract, 0.24 to 3.07 mg RU/g crude extract, and 1.31 to 7.89 mg EE/g crude extract, respectively (Table 3). The bark methanolic extract contained the highest TPC and TFC, and the second highest TSC ($p < 0.05$) (Fig. 4). A two-way ANOVA revealed a significant effect of plant parts and solvents on TPC ($F = 51.830$, $p < 0.001$ for plant parts; $F = 209.899$, $p < 0.001$ for solvents) and TFC ($F = 80.876$, $p < 0.001$ for plant parts; $F = 160.703$, $p < 0.001$ for solvents). Notably, the interaction between plant parts and solvents also had a significant effect ($F = 50.424$, $p < 0.001$) on TPC (Fig. 4A) and TFC ($F = 126.575$, $p < 0.001$) (Fig. 4B). The analysis showed similar effects on TSC of plant parts ($F = 73.781$, $p < 0.001$) and solvents ($F = 24.045$, $p < 0.001$) and their interactions ($F = 4.484$, $p = 0.001$) but the effects were less significant (Fig. 4C).




Figure 4: Interaction effects of plant parts and solvents on (A) total phenolic content (TPC), (B) total flavonoid content (TFC), and (C) total saponin content (TSC) in *C. inophyllum* extracts. Different lowercase letters denote significant differences ($p < 0.05$) among means, $n = 3$; error bars represent \pm SD. EtOAc: ethyl acetate, MeOH: methanol.

The differences in TPC, TFC, and TSC across the extracts could be influenced by environmental factors, the chemical composition of the plant material, soil conditions, harvest maturity, and post-harvest storage practices [38,39]. In the present study, the effects of the different solvents were explored. The methanolic extracts yielded the highest overall TPC and TFC, whereas the hexane extracts yielded the highest TSC (Fig. 5). When averaged across solvents, the twig and bark extracts contained the highest TPCs among the plant parts (0.83 ± 0.52 mg GAE/g CE for twig and 0.91 ± 0.68 mg GAE/g CE for bark). The leaf and twig extracts yielded the highest TFCs (1.91 ± 0.20 mg RU/g CE for leaf and 1.81 ± 0.86 mg RU/g CE for twig). The bark extract yielded the highest TSC (6.91 ± 1.25 mg EE/g CE). Among solvents (averaged across plant parts), the TPC and TFC were highest in the methanolic extracts (0.97 ± 0.62 mg GAE/g CE for TPC and 1.88 ± 0.98 mg RU/g CE for TFC), whereas TSC was highest in hexane extracts (4.96 ± 2.27 mg EE/g CE) (Fig. 5, Tables S1 and S2).




Figure 5: Total phenolic content (TPC), total flavonoid content (TFC), and total saponin content (TSC) were averaged across solvents and across plant parts in *C. inophyllum* extracts. Different lowercase letters denote significant differences ($p < 0.05$) among means; error bars represent \pm SD. $n = 15$ for solvents $n = 9$ for plant part EtOAc: ethyl acetate, MeOH: methanol.

The results indicated that the solvent used determined whether phenolics, flavonoids, or saponins were the major phytochemicals extracted. This finding is consistent with prior research that highlighted the influence of solvent polarity on the solubility of bioactive compounds [19]. Methanol is a polar solvent that is particularly effective in extracting phenolics and flavonoids, while hexane, a non-polar solvent, is more effective in extracting saponins. These results were consistent with the principles of solvent extraction, where the choice of solvent is crucial for optimizing yield based on the chemical nature of target compounds [40]. An understanding of the distribution of bioactive compounds among the various plant parts is also crucial. Twig and bark material yield the highest phenolic contents, while leaf and twig yield

the highest flavonoid contents. This differential distribution could be attributed to the adaptive mechanisms of plants and the different roles different tissues play in protecting against environmental stressors such as UV radiation [41] and pathogens [42].

3.4 Determination of Antioxidant Activity

The antioxidant activity of *C. inophyllum* extracts was evaluated using the DPPH and ABTS methods. Bioactive compounds such as phenolics, saponins, terpenoids, and anthraquinones, particularly those containing hydroxyl groups (–OH), act as scavengers, neutralizing lone-pair electrons in the reaction systems, causing visible color changes. DPPH changes from purple to colorless, while ABTS changes from light blue to yellow. Table 3 shows the antioxidant activities of *C. inophyllum* extracts from flower, fruit, leaf, twig, and bark extracted using different solvents. The methanolic bark extract demonstrated the highest antioxidant activity (0.004 ± 0.000 mg/L), which was comparable to the antioxidant activity of Vitamin C, but better than that of the synthetic antioxidant BHT (0.011 ± 0.000 mg/L). This strong antioxidant activity was likely due to the high content of natural phenolic and flavonoid compounds in the methanolic extract. The presence of hydroxyl groups in phenolic compounds plays a crucial role in trapping and neutralizing free radicals [7,43]. Therefore, this effectiveness is likely related to the high polarity of compounds extracted with methanol [9].

The Pearson correlation coefficient was used to study the relationship between phytochemical contents in *C. inophyllum* extracts and antioxidant activities (Fig. 6). A strong positive correlation was observed between TPC and TFC ($r = 0.776$, $p < 0.001$). However, antioxidant activity was significantly correlated only with TPC ($r = -0.559$, $p < 0.001$ for DPPH; $r = -0.484$, $p = 0.001$ for ABTS), and not with TFC ($r = -0.072$, $p = 0.639$ for DPPH; $r = -0.073$, $p = 0.633$ for ABTS) or TSC ($r = 0.038$, $p = 0.802$ for DPPH; $r = -0.081$, $p = 0.599$ for ABTS). The differences in antioxidant activity could be attributed to the specific types of phenolic, flavonoid, and saponin compounds present in each extract or to the presence of other primary metabolites, such as sugars, carbohydrates, proteins, and amino acids [19,44]. Similarly, Vittaya et al. [14] reported a positive correlation ($r = 0.876$, $p = 0.002$) between phenolic and flavonoid contents and antioxidant activity in *Ampelocissus martini* extracts.




Figure 6: Total phenolic content (TPC), total flavonoid content (TFC), and total saponin content (TSC) in each plant part were averaged across solvents. TPC and TFC were positively correlated with each other but negatively correlated with antioxidant activity [2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azinobis-3-ethylbenzothiazole-6-sulfonic acid (ABTS)] of flower, fruit, leaf, twig, and bark of *C. inophyllum*. The data are presented as means \pm SD. Pearson's correlation coefficient (r) between each pair of plant parts and phytochemical contents ($n = 9$), ** $p < 0.001$.

3.5 Determination of Antibacterial Activity

All extracts of *C. inophyllum* showed antibacterial activity against the seven tested bacterial strains, which included three Gram positive and four Gram negative species. The diameters of the inhibition zones for Gram positive bacteria ranged from 10.57–17.16 mm against *B. cereus*, 8.28–17.96 mm against *S. aureus*, and 8.11–9.61 mm against *S. epidermidis*. For Gram negative bacteria, the diameters of the inhibition zones ranged from 7.57–9.08 mm against *E. coli*, 8.27–17.15 mm against *S. typhi*, 6.32–8.44 mm against *K. pneu*, and 8.53–18.78 mm against *P. aeruginosa* (Table 4). Moreover, the antibacterial effects varied depending on the plant parts, the solvents used for extraction, and their combinations. For *B. cereus*, *S. aureus*, and *S. typhi*, plant parts, solvents, and the interaction between plant parts and solvents were highly significant ($p < 0.001$). The best antibacterial activities were found in the hexane extracts from bark and twig. For *S. epidermidis*, plant parts and solvents were significant ($p = 0.01$), though the interaction between plant parts and solvents was not as strong. The methanolic bark extract demonstrated the strongest antibacterial activity against this bacterium. For *E. coli* and *K. pneumoniae*, plant parts were significant ($p = 0.002$ and $p < 0.001$, respectively), and the interaction between plant parts and solvents was also significant ($p = 0.007$ and $p < 0.001$). The methanolic extracts of twig and bark showed the highest activity against *E. coli*, while the hexane flower extract was most effective against *K. pneumoniae*. In the case of *P. aeruginosa*, both plant parts and solvents were significant ($p < 0.001$), along with their interaction ($p = 0.01$). The best antibacterial activities were found in the hexane extract of fruit. Overall, these results demonstrated that the hexane and methanolic extracts from bark were more effective against Gram positive bacteria than Gram negative bacteria. However, hexane extracts from twig, flower, and fruit were notably more effective against Gram negative bacteria.

Table 4: Zones of inhibition from methanolic extracts of *C. inophyllum* against seven pathogenic bacteria. Two-way analysis of variance showed the effects of plant parts (PP), solvent used (S), and their interaction.

Parts	Solvents	Zone of Inhibition Diameter (mm)													
		Gram Positive						Gram Negative							
		<i>B. cereus</i>		<i>S. aureus</i>		<i>S. epidermidis</i>		<i>E. coli</i>		<i>S. typhi</i>		<i>K. pneu</i>		<i>P. aeruginosa</i>	
Flower*	H	15.07 ± 1.03 ^{c-e,AB}	14.95 ± 0.29 ^{de,AB}	8.48 ± 0.51 ^{b-d,C}	8.54 ± 0.51 ^{bc,C}	16.63 ± 0.79 ^{c,A}	8.44 ± 0.31 ^{b,C}	13.92 ± 3.18 ^{d-f,B}							
	E	14.46 ± 0.21 ^{de,A}	14.81 ± 0.85 ^{de,A}	8.61 ± 0.58 ^{b-d,C}	8.25 ± 0.60 ^{c-f,C}	14.35 ± 0.94 ^{ef,A}	7.70 ± 0.39 ^{cd,C}	12.01 ± 2.74 ^{f,B}							
	M	14.09 ± 0.75 ^{de,B}	13.66 ± 0.22 ^{ef,B}	8.58 ± 0.30 ^{b-d,C}	7.68 ± 0.65 ^{d-f,CD}	16.62 ± 0.35 ^{c,A}	6.85 ± 0.25 ^{e-g,D}	12.55 ± 2.18 ^{ef,B}							
Fruit	H	15.32 ± 1.55 ^{b-d,B}	15.71 ± 1.53 ^{cd,B}	9.10 ± 1.40 ^{b-d,C}	8.12 ± 0.10 ^{c-f,C}	17.04 ± 1.23 ^{bc,AB}	8.11 ± 0.07 ^{bc,C}	18.78 ± 0.27 ^{b,A}							
	E	14.27 ± 1.52 ^{de,B}	14.15 ± 0.60 ^{ef,B}	8.60 ± 0.52 ^{b-d,C}	7.45 ± 0.25 ^{f,C}	17.03 ± 0.80 ^{bc,A}	7.89 ± 0.47 ^{b-d,C}	17.07 ± 0.99 ^{bc,A}							
	M	13.30 ± 0.64 ^{d-f,B}	14.11 ± 0.77 ^{ef,B}	8.93 ± 0.83 ^{b-d,C}	8.37 ± 0.22 ^{b-e,CD}	16.73 ± 1.22 ^{c,A}	7.33 ± 0.46 ^{d-f,D}	15.86 ± 1.05 ^{cd,A}							
Leaf	H	10.57 ± 0.06 ^{h,A}	8.28 ± 0.34 ^{h,BC}	8.11 ± 0.22 ^{de,BC}	8.04 ± 0.36 ^{c-f,BC}	8.50 ± 0.46 ^{h,B}	7.80 ± 0.23 ^{b-d,C}	8.53 ± 0.39 ^{g,B}							
	E	11.11 ± 1.02 ^{gh,A}	8.61 ± 0.39 ^{h,BC}	7.31 ± 0.54 ^{e,D}	7.74 ± 0.61 ^{c-f,CD}	8.27 ± 0.32 ^{h,BCD}	7.63 ± 0.59 ^{cd,CD}	9.03 ± 0.60 ^{g,B}							
	M	11.73 ± 2.47 ^{f-h,A}	8.92 ± 0.48 ^{gh,BC}	8.17 ± 0.38 ^{de,BC}	7.57 ± 0.51 ^{ef,C}	8.72 ± 0.81 ^{h,CD}	7.66 ± 0.57 ^{cd,C}	9.69 ± 0.37 ^{g,B}							
Twig	H	17.04 ± 1.40 ^{bc,A}	16.77 ± 1.05 ^{bc,A}	8.79 ± 1.06 ^{b-d,C}	8.13 ± 0.49 ^{c-f,C}	18.47 ± 1.46 ^{b,A}	6.32 ± 0.07 ^{g,D}	14.90 ± 0.67 ^{c-e,B}							
	E	10.57 ± 0.85 ^{h,AB}	10.19 ± 1.47 ^{g,B}	8.01 ± 0.39 ^{de,C}	8.42 ± 0.65 ^{b-e,C}	11.78 ± 0.51 ^{g,A}	6.32 ± 0.12 ^{g,D}	9.26 ± 0.23 ^{g,BC}							
	M	13.05 ± 1.70 ^{e-g,B}	12.92 ± 0.31 ^{f,B}	9.37 ± 0.27 ^{bc,C}	9.08 ± 0.23 ^{b,C}	15.88 ± 0.54 ^{cd,A}	7.44 ± 0.57 ^{c-e,D}	13.07 ± 0.65 ^{ef,B}							
Bark	H	17.16 ± 0.96 ^{b,AB}	17.96 ± 0.71 ^{b,A}	8.31 ± 0.45 ^{c-e,C}	8.18 ± 0.02 ^{cb-d,C}	17.15 ± 0.49 ^{bc,AB}	6.88 ± 0.30 ^{e-g,D}	16.42 ± 0.63 ^{c,B}							
	E	13.77 ± 0.47 ^{d-f,C}	15.95 ± 0.56 ^{cd,A}	8.44 ± 0.06 ^{c-e,D}	8.24 ± 0.20 ^{c-f,D}	14.91 ± 1.03 ^{de,B}	6.70 ± 0.31 ^{fg,E}	13.63 ± 0.30 ^{d-f,C}							
	M	13.15 ± 0.43 ^{d-g,A}	13.84 ± 0.97 ^{ef,A}	9.61 ± 0.28 ^{b,A}	9.08 ± 0.23 ^{b,B}	13.22 ± 1.35 ^{fg,A}	7.27 ± 0.16 ^{d-f,C}	13.29 ± 0.68 ^{ef,A}							
Gentamycin		23.15 ± 0.12 ^{a,C}	23.91 ± 0.06 ^{a,B}	22.01 ± 10.34 ^{a,D}	20.74 ± 0.04 ^{a,E}	26.14 ± 0.04 ^{a,A}	18.69 ± 0.15 ^{a,F}	23.41 ± 0.17 ^{a,C}							
Two-Way ANOVA															
Variable	df	F	P	F	P	F	P	F	P	F	P	F	P		
PP	4	13.960	<0.001	110.964	<0.001	3.944	0.011	5.260	0.002	125.538	<0.001	16.490	<0.001	44.441	<0.001
S	2	15.732	<0.001	31.289	<0.001	5.328	0.010	2.300	0.118	24.618	<0.001	2.106	0.139	11.691	<0.001
PP × S	8	5.198	<0.001	11.319	<0.001	1.173	0.347	3.389	0.007	9.759	<0.001	6.724	<0.001	3.1192	0.010
Error	30														

*Data from the flower extracts are reused in this article by permission from Elsevier (License No. 5795780615723) [10]. Zones of inhibition expressed as means (mm) ± SD ($n = 3$). Means ± SD in a column superscripted with lowercase letters and in a row with uppercase letters denote significant differences ($p < 0.05$). H: hexane, E: ethyl acetate, and M: methanol.

The stronger activity demonstrated by flower and twig extracts against *S. typhi* compared to other bacteria was consistent with the findings of previous research [10]. Also, the fruit extract was active against both *S. typhi* and *P. aeruginosa*. The leaf extract exhibited the highest activity against *B. cereus* compared to other plant parts, while the bark extract was most effective against *S. aureus*. Moreover, the chemical compositions in the flower extracts were rich in sesquiterpenoids, triterpenoids, fatty acids, and fatty acid derivatives, therefore they could inhibit the growth of Gram positive and Gram negative bacteria, especially *S. typhi* [10].

The Minimum Inhibitory Concentration (MIC) of the *C. inophyllum* extracts ranged from 0.098 to >50 µg/mL against Gram positive bacteria and from 0.19 to >100 µg/mL against Gram negative bacteria (Table 5). All extracts showed strong activity against *B. cereus* with low MIC values between 0.098 µg/mL to 0.78 µg/mL and decreasing activity against *S. typhi*, *S. aureus*, *P. aeruginosa*, *S. epidermidis*, and *E. coli*, and the least activity against *K. pneumoniae*. The antibacterial mechanism is impacted by the molecular weight of compounds, which influences their diffusion through agar, and therefore the chemical composition of the extracts contributed to the variations in sensitivity observed between Gram positive and Gram negative bacteria [45,46].

Table 5: Minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC) of *C. inophyllum* extracts.

Parts	Solvents	MIC/MBC (µg/mL)													
		Gram Positive								Gram Negative					
		<i>B. cereus</i>		<i>S. aureus</i>		<i>S. epidermidis</i>		<i>E. coli</i>		<i>S. typhi</i>		<i>K. pneumoniae</i>		<i>P. aeruginosa</i>	
		MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Flower*	H	<0.098	3.12	12.5	50	12.5	25	25	>50	12.5	>100	50	>100	25	>100
	E	0.39	25	6.25	>50	6.25	25	6.25	50	12.5	>100	12.5	100	6.25	100
	M	0.098	3.12	6.25	6.25	6.25	50	12.5	>50	12.5	>100	25	>100	12.5	>100
Fruit	H	<0.098	1.51	25	50	12.5	>50	12.5	>50	12.5	>100	25	100	12.5	>100
	E	<0.098	1.51	3.12	25	12.5	>50	25	>50	12.5	>100	25	100	12.5	100
	M	<0.098	1.51	12.5	25	12.5	>50	12.5	>50	6.25	>100	25	100	6.25	>100
Leaf	H	0.39	50	6.25	50	12.5	>50	25	>50	6.25	>100	50	>100	25	>100
	E	0.78	50	3.12	>50	12.5	>50	12.5	>50	3.12	>100	25	>100	12.5	>100
	M	0.39	50	3.12	50	12.5	>50	6.25	>50	1.51	100	25	>100	3.12	100
Twig	H	<0.098	1.51	50	50	12.5	>50	25	50	0.19	50	50	>100	12.5	>100
	E	0.098	25	6.25	50	12.5	>50	12.5	>50	1.51	100	25	>100	6.25	>100
	M	0.098	6.25	6.25	12.5	6.25	>50	12.5	>50	0.78	50	12.5	>100	1.51	100
Bark	H	<0.098	0.78	25	50	12.5	>50	25	>50	12.5	>100	50	>100	12.5	>100
	E	<0.098	0.19	12.5	50	12.5	>50	12.5	>50	6.25	>100	25	>100	12.5	>100
	M	0.098	0.19	6.25	6.25	6.25	>50	12.5	50	1.51	100	12.5	100	0.39	50

*Data from the flower extracts are reused in this article by permission from Elsevier (License No. 5795780615723) [10]. Values are presented as mean results from triplicate determinations ($n = 3$). H: hexane, E: ethyl acetate, and M: methanol.

Fig. 7 presents the heatmap of Pearson correlations between TPC, TPC, and TSC and antibacterial activity. The color gradient indicates the strength of the correlation (ranging from -1 to 1). Negative correlations are blue and positive correlations red. The results show that TPC was significantly positively correlated with antibacterial activity against *S. epidermidis* and *E. coli*. However, TFC displayed a highly significant negative correlation with antibacterial activity against *B. cereus*, *S. aureus*, *E. coli*, *K. pneumoniae*, and *P. aeruginosa*. Additionally, TFC showed a significant positive correlation with activity against *E. coli* but a significant negative correlation with activity against *S. typhi*. As mentioned above, these variations in sensitivity were associated with the chemical composition of the extracts.




Figure 7: Correlation heatmap showing the relationship between phytochemical contents and antibacterial activities of *C. inophyllum* extracts. Asterisks (*, **) indicate significance at $p < 0.05$ and $p < 0.001$, respectively.

3.6 Principal Component Analysis (PCA) of Phytochemicals and Biological Activities in *C. inophyllum* Extracts

The first two principal components, PC1 and PC2, accounted for 36.3% and 25.4% of the total variance, respectively, and together explained most of the variation observed in the dataset. The loadings of variables on PC1 and PC2, along with the scores for each plant part-solvent combination, are presented in a PCA biplot (Fig. 8). The arrows indicate the direction and strength of correlation for each variable, with longer arrows representing higher contributions to the principal components. Samples located close to a particular variable arrow represent extracts with higher values for that attribute.

The first component PC1, showed that positive scores were correlated with higher antibacterial activity against *B. cereus*, *S. aureus*, *S. epidermidis*, *S. typhi*, and *P. aeruginosa*. This finding indicated that hexane extracts of bark and twig possessed strong antibacterial activity. The stronger antibacterial efficacy observed in non-polar hexane extracts could be attributed to the presence of lipophilic compounds, such as terpenoids and fatty acids, which are known to disrupt bacterial cell membranes and interfere with lipid-dependent metabolic processes [22,46]. In contrast, negative PC1 scores were associated with lower antibacterial activity and the findings suggested that leaf extracts, which were particularly flavonoid-rich, did not exhibit strong antibacterial activity against the tested bacterial strains. This observation was consistent with previous studies, which have shown that the antibacterial efficacy of specific flavonoid structures can vary depending on the bacterial species and extraction methods [43]. The second component PC2, indicated that positive scores were associated with higher TPC, TFC, and antibacterial activity against *E. coli* and *S. epidermidis*. Methanol extracts of bark and twigs exhibited these properties. Conversely, negative PC2 scores corresponded to lower IC₅₀ values for DPPH and ABTS assays, indicating higher antioxidant activities, which were observed in ethyl acetate and methanolic leaf extracts, and lower antibacterial activity, especially against *K. pneumoniae*, notably in fruit and leaf hexane extracts. The inverse relationship between antioxidant and antibacterial activities could be attributed to the different roles that bioactive compounds serve in each function. Antioxidants, such as phenolic acids and flavonoids, are primarily involved in neutralizing free radicals and protecting cells from oxidative damage [47,48]. Their mechanism of action is

based on electron donation or radical scavenging, which may not directly impact bacterial cell walls or metabolic pathways [49,50].




Figure 8: Principal Component Analysis (PCA) biplot illustrating the relationships between phytochemical contents (TPC, TFC, and TSC), antioxidant activities (DPPH and ABTS), antibacterial activities against various bacterial strains, and the different solvents used for extracting aerial parts of *C. inophyllum*. B: bark, Fr: fruit, Fw: flower, L: leaf, T: twig. Hex: hexane, EtOAc: ethyl acetate, MeOH: methanol. BC: *B. cereus*, SA: *S. aureus*, SE: *S. epidermidis*, EC: *E. coli*, ST: *S. typhi*, KP: *K. pneumoniae*.

3.7 Cytotoxicity and Anticancer Activity

The anticancer activities of *C. inophyllum* extracts are shown in Table 6. The most significant anticancer activity was observed in the leaf and bark extracts, especially those extracted with ethyl acetate and hexane. The ethyl acetate leaf extract exhibited the strongest effect against NCI-H187, with an IC_{50} value of 10.09 $\mu\text{g/mL}$, followed by the hexane extract, with an IC_{50} value of 17.45 $\mu\text{g/mL}$. These results suggest that solvents with intermediate polarity (ethyl acetate) and non-polar solvents (hexane) effectively extract the bioactive compounds responsible for anticancer activity, particularly from the leaf of *C. inophyllum*, which produced high TPC and TFC (Table 3). The anticancer activities of phenolics and flavonoids are well known, and the high TPC and TFC of ethyl acetate and hexane leaf extracts correlate with the strong anticancer activity observed in these extracts [51]. The ethyl acetate and hexane bark extracts showed anticancer activity, with IC_{50} values of 19.80 $\mu\text{g/mL}$ and 18.03 $\mu\text{g/mL}$, respectively, further supporting the presence of active bioingredients and also significant saponin content in comparison to other parts. Saponins may have contributed to the observed anticancer effects, as they are known to inhibit cancer cell proliferation and induce apoptosis [29].

Furthermore, certain extracts demonstrated selective cytotoxicity against cancer cells, while also showing moderate cytotoxicity against Vero cells. For instance, the leaf and bark extracts exhibited moderate toxicity (65.58% for hexane leaf extract and 84.88% for ethyl acetate leaf extract). The cytotoxicity of fruit, twig, and bark extracts also exceeded the 50% threshold for cytotoxicity. While the flower extracts were non-cytotoxic to both cancerous and normal cells, their phytochemical composition (particularly their lower TPC and TFC) likely accounts for the lack of anticancer activity. Previous studies have reported that ethanolic extracts of *C. inophyllum* contained high phenolic and flavonoid contents [8], and other anticancer-related

compounds such as curcumin, resveratrol, epigallocatechin gallate, and thymoquinone [52–55]. Further support for the anticancer potential of this plant was recently provided by Ruangsuriya et al. [56], who found the three most abundant phytochemicals in *C. inophyllum* extracts to be 5-hydroxy methylfurfural (5-HMF), antiarol, and syringol, which are classified as furans and phenols. However, despite the significant activity observed in the present study against NCI-H187, no notable effects were observed against the KB and MCF7 cancer cell lines across all extracts. This lack of activity suggests that the bioactive compounds in *C. inophyllum* may not specifically target the pathways relevant to these cancer types, or that higher concentrations may be required to observe activity. These findings highlight the potential of this plant in the use of traditional medicine for cancer therapy. Nonetheless, the higher cytotoxicity observed in some extracts, particularly those from bark and twig, warrants caution and highlights the need for further investigation to balance anticancer efficacy with safety.

Table 6: Cytotoxicity of *C. inophyllum* extracts against three cancer cell lines: oral cavity cancer (KB), breast cancer (MCF7), small cell lung cancer (NCI-H187), along with IC₅₀, and cytotoxicity against African green monkey kidney (Vero) cells.

Parts	Solvents	KB	MCF7	NCI-H187	IC ₅₀ (μg/mL)	Vero Cell (% Cytotoxicity)
Flower	H	Inactive	Inactive	Inactive	-	Non-cytotoxic
	E	Inactive	Inactive	Inactive	-	Non-cytotoxic
	M	Inactive	Inactive	Inactive	-	Non-cytotoxic
Fruit	H	Inactive	Inactive	Inactive	-	Non-cytotoxic
	E	Inactive	Inactive	56.89%	48.51 ± 0.28	81.60%
	M	Inactive	Inactive	Inactive	-	Non-cytotoxic
Leaf	H	Inactive	Inactive	99.11%	17.45 ± 1.51	65.58%
	E	Inactive	Inactive	93.29%	10.09 ± 1.43	84.88%
	M	Inactive	Inactive	Inactive	-	Non-cytotoxic
Twig	H	Inactive	Inactive	Inactive	-	Non-cytotoxic
	E	Inactive	Inactive	91.12%	18.70 ± 0.27	91.98%
	M	Inactive	Inactive	Inactive	-	Non-cytotoxic
Bark	H	Inactive	Inactive	82.42%	18.03 ± 0.09	92.08%
	E	Inactive	Inactive	83.96%	19.80 ± 0.92	87.05%
	M	Inactive	Inactive	Inactive	-	Non-cytotoxic
Doxorubicin		0.90 ± 0.12	9.38 ± 0.23	0.09 ± 0.02	-	-
Ellipticine		2.49 ± 0.20	-	2.15 ± 0.72	-	-
Tamoxifen		-	7.95 ± 0.84	-	-	-

Note: % Cytotoxicity: <50% interpreted non-cytotoxic, ≥50% interpreted cytotoxic. H: hexane, E: ethyl acetate, and M: methanol.

3.8 In Silico ADME Analysis of Selected Phytochemical Compounds from *C. inophyllum* Extract

The bioactive compounds found by Vittaya et al. [10] in a flower extract of *C. inophyllum* (Fig. 9) were studied by *in silico* ADME analysis. Selected phytochemical compounds from *C. inophyllum* flower extract revealed diverse pharmacokinetic profiles, highlighting their potential as drug candidates (Table 7 and Table S3). The analysis revealed that eugenol, caryophyllene oxide, farnesol, and palmitic acid demonstrated good gastrointestinal absorption, supporting their potential bioavailability if taken orally [57–60]. Other compounds, including α-copaene and β-amyrin, displayed lower GI absorption and may require formulation optimization to enhance their bioavailability. Additionally, blood-brain barrier permeability was predicted for eugenol and caryophyllene oxide, indicating potential for treating central nervous system disorders. These compounds have demonstrated neuroprotective and biological activities in previous studies [61–63]. Further analysis identified various cytochrome P450 (CYP) enzymes, with compounds like α-copaene and palmitic acid identified as inhibitors of CYP1A2 and CYP2C9, emphasizing the importance of considering the potential risks of drug–drug interactions [64,65]. Most compounds adhered to Lipinski's Rule of Five, indicating favorable drug-likeness and suitability for further development [66]. Collectively, these results

indicated that *C. inophyllum* extracts possess significant antioxidant and antibacterial activities for potential applications in food and cosmetics, while ADME analysis supports the therapeutic potential of selected phytochemicals as candidates for further exploration in pharmaceutical development.




Figure 9: Selected phytochemical compounds identified in a *C. inophyllum* flower extract.

Table 7: *In Silico* ADME analysis of physicochemical and pharmacokinetic properties of selected phytochemical compounds identified in a *C. inophyllum* flower extract.

Compounds	Formula	Mol. Weight (g/mol)	Num. Heavy Atoms	Num. Rotatable Bonds	TPSA (Å ²)	Consensus LogP o/w	Water Solubility Class	GI Absorption	BBB Permeation	P-gp Substrate	CYP Inhibitors	Druglikeness Lipinski	Bioavailability Score	PAINS	Synthetic Accessibility Score
Eugenol	C ₁₀ H ₁₂ O ₂	164.20	12	3	29.46	2.25	Soluble	High	Yes	No	CYP1A2	Yes (0 violation)	0.55	0	1.58
α-Copaene	C ₁₅ H ₂₄	204.35	15	1	0.00	4.30	Soluble	Low	Yes	No	CYP1A2, CYP2C19, CYP2C9	Yes (1 violation)	0.55	0	4.62
β-Caryophyllene	C ₁₅ H ₂₄	204.35	15	0	0.00	4.24	Soluble	Low	No	No	CYP2C19, CYP2C9	Yes (1 violation)	0.55	0	4.51
α-Murolene	C ₁₅ H ₂₄	204.35	15	1	0.00	4.08	Soluble	Low	No	No	CYP2C19, CYP2C9	Yes (1 violation)	0.55	0	4.35
γ-Cadinene	C ₁₅ H ₂₄	204.35	15	1	0.00	4.18	Soluble	Low	No	No	CYP2C19, CYP2C9	Yes (1 violation)	0.55	0	4.35
δ-Cadinene	C ₁₅ H ₂₄	204.35	15	1	0.00	4.12	Soluble	Low	No	No	CYP2C19, CYP2C9	Yes (1 violation)	0.55	0	4.14
Caryophyllene Oxide	C ₁₅ H ₂₄ O	220.35	16	0	12.53	3.68	Soluble	High	Yes	No	CYP2C19, CYP2C9	Yes (0 violation)	0.55	0	4.35
Farnesol	C ₁₅ H ₂₆ O	222.37	16	7	20.23	3.68	Soluble	High	Yes	No	CYP2C19, CYP2C9	Yes (0 violation)	0.55	0	4.35
Palmitic Acid	C ₁₆ H ₃₂ O ₂	256.42	18	14	37.30	5.20	Moderately soluble	High	Yes	No	CYP1A2, CYP2C9	Yes (1 violation)	0.85	0	2.31
Phytol	C ₂₀ H ₄₀ O	296.53	21	13	20.23	6.25	Moderately soluble	Low	No	Yes	CYP2C9	Yes (1 violation)	0.55	0	4.30
α-Linolenic Acid	C ₁₈ H ₃₀ O ₂	278.43	20	13	37.30	6.25	Moderately soluble	Low	No	Yes	CYP2C9	Yes (1 violation)	0.55	0	4.30
2-cis-Geranylgeraniol	C ₂₀ H ₃₄ O	290.48	21	10	20.23	6.25	Moderately soluble	Low	No	Yes	CYP2C9	Yes (1 violation)	0.55	0	4.30
β-Amyrin	C ₃₀ H ₅₀ O	426.72	31	0	20.23	7.18	Poorly soluble	Low	No	No	None	Yes (1 violation)	0.55	0	6.04

Note: TPSA: Topological polar surface area, LogP o/w: n-octanol/water partition coefficient (consensus log Po/w value is the arithmetic means of the five predictive values), GI: Gastrointestinal, BBB: Blood-brain barrier, P-gp: P glycoprotein, CPY: Cytochrome P450, Lipinski: Lipinski (Pfizer) filter, PAINS: Pan assay interference, Yes (0 violation): meet all criteria, Yes (1 violation): MLOGP > 4.15.

4 Conclusion

C. inophyllum flower, fruit, leaf, twig, and bark were extracted by maceration in hexane, ethyl acetate, and methanol. The effects of the different plant parts and solvents on the phytochemical profiles of the extracts were investigated, including the effects of interactions between plant parts and solvents on biological activities. The extraction yield of *C. inophyllum* varied based on the solvent used, with methanol yielding the most extract, followed by ethyl acetate and hexane. Total phenolic and flavonoid contents were significantly correlated. The highest total phenolic and flavonoid contents were observed in ethyl acetate and methanolic extracts, which exhibited the highest antioxidant activity. The highest total saponin content was found in hexane extracts, except for the fruit extract. All extracts demonstrated inhibitory effects against Gram-positive and negative bacteria, with a particularly strong effect against *S. typhi*. Additionally, the leaf and bark extracts showed promising anticancer activity, especially against the NCI-H187 lung cancer cell line, suggesting potential for therapeutic application. However, further research is needed to assess its broader anticancer efficacy and safety profile. SwissADME screening identified eugenol and caryophyllene oxide as promising bioactive candidates with good gastrointestinal absorption, blood–brain barrier permeability, favorable topological polar surface areas, and minimal cytochrome P450 enzyme interactions. These characteristics suggest potential suitability for both systemic and central nervous system applications. Farnesol and palmitic acid also showed good oral bioavailability potential. Although the pharmacokinetic predictions were limited to compounds identified in the flower extracts, these findings provide a valuable foundation for further investigation. A more comprehensive identification and pharmacokinetic evaluation of compounds from other plant parts is needed to fully explore their therapeutic potential. Further studies incorporating complete ADMET profiling will also be essential to better clarify the safety and therapeutic relevance of *C. inophyllum* metabolites. Overall, these results highlight the potential of *C. inophyllum* extracts for applications in the food, cosmetic, and pharmaceutical industries.

Acknowledgement: This project was financially supported by the Faculty of Science and Fisheries Technology, Rajamangala University of Technology Srivijaya, Trang campus. The authors would like to thank the Department of National Parks, Wildlife and Plant Conservation, Thailand for the specimen voucher of *C. inophyllum* and the National Center for Genetic Engineering and Biotechnology (BIOTEC), Thailand for pathogenic bacteria. Special thanks go to Thomas Duncan Coyne for proofing and editing the English text of the manuscript.

Funding Statement: The authors received no specific funding for this study.

Author Contributions: The authors confirm contribution to the paper as follows: Conceptualization, Luksamee Vittaya; Formal analysis, Luksamee Vittaya; Investigation, Luksamee Vittaya, Chakhriya Chalad and Sittichoke Janyong; Methodology, Luksamee Vittaya and Chakhriya Chalad; Supervision, Luksamee Vittaya; Validation, Luksamee Vittaya and Sittichoke Janyong; Visualization, Luksamee Vittaya and Nararak Leesakul; Writing—original draft, Luksamee Vittaya; Writing—review & editing, Nararak Leesakul. All authors reviewed the results and approved the final version of the manuscript.

Availability of Data and Materials: The authors confirm that the data supporting the findings of this study are available within the article and its Supplementary Materials.

Ethics Approval: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest to report regarding the present study.

Supplementary Materials: The supplementary material is available online at <https://www.techscience.com/doi/10.32604/phyton.2025.074891/s1>.

References

- Watts N, Amann M, Arnell N, Ayeb-Karlsson S, Beagley J, Belesova K, et al. The 2020 report of the Lancet countdown on health and climate change: responding to converging crises. *Lancet*. 2021;397(10269):129–70. [[CrossRef](#)].
- Guzmán-Trampe S, Ceapa CD, Manzo-Ruiz M, Sánchez S. Synthetic biology era: improving antibiotic's world. *Biochem Pharmacol*. 2017;134:99–113. [[CrossRef](#)].
- Olatunji OJ, Olatunde OO, Jayeoye TJ, Singh S, Nalinbenjapun S, Sripetthong S, et al. New insights on *Acanthus ebracteatus* Vahl: UPLC-ESI-QTOF-MS profile, antioxidant, antimicrobial and anticancer activities. *Molecules*. 2022;27(6):1981. [[CrossRef](#)].
- Chan EWC, Ng YK, Wong SK, Chan HT. *Pluchea indica*: an updated review of its botany, uses, bioactive compounds and pharmacological properties. *Pharm Sci Asia*. 2022;49(1):77–85. [[CrossRef](#)].
- Ferdosh S. The extraction of bioactive agents from *Calophyllum inophyllum* L., and their pharmacological properties. *Sci Pharm*. 2024;92(1):6. [[CrossRef](#)].
- Vittaya L, Charoendat U, Janyong S, Ui-eng J, Leesakul N. Comparative analyses of saponin, phenolic, and flavonoid contents in various parts of *Rhizophora mucronata* and *Rhizophora apiculata* and their growth inhibition of aquatic pathogenic bacteria. *J App Pharm Sci*. 2022;12(11):111–21. [[CrossRef](#)].
- Audah KA, Ettin J, Darmadi J, Azizah NN, Anisa AS, Hermawan TDF, et al. Indonesian mangrove *Sonneratia caseolaris* leaves ethanol extract is a potential super antioxidant and anti methicillin-resistant *Staphylococcus aureus* drug. *Molecules*. 2022;27(23):8369. [[CrossRef](#)].
- Cassien M, Mercier A, Thétiot-Laurent S, Culcasi M, Riquebourg E, Asteian A, et al. Improving the antioxidant properties of *Calophyllum inophyllum* seed oil from French Polynesia: development and biological applications of resinous ethanol-soluble extracts. *Antioxidants*. 2021;10(2):199. [[CrossRef](#)].
- Hapsari S, Yohed I, Kristianita RA, Jadid N, Aparamarta HW, Gunawan S. Phenolic and flavonoid compounds extraction from *Calophyllum inophyllum* leaves. *Arab J Chem*. 2022;15(3):103666. [[CrossRef](#)].
- Vittaya L, Chalad C, Ratsameepakai W, Leesakul N. Phytochemical characterization of bioactive compounds extracted with different solvents from *Calophyllum inophyllum* flowers and activity against pathogenic bacteria. *S Afr N J Bot*. 2023;154:346–55. [[CrossRef](#)].
- Ojah EO, Moronkola DO, Petrelli R, Nzekoue FK, Cappellacci L, Giordani C, et al. Chemical composition of ten essential oils from *Calophyllum inophyllum* Linn and their toxicity against *Artemia Salina*. *Eur J Pharm Med Res*. 2019;6(12):185–94.
- Ojah EO, Moronkola DO, Osamudiamen PM. Antioxidant assessment of characterised essential oils from *Calophyllum inophyllum* Linn using 2, 2-diphenyl-1-picrylhydrazyl and hydrogen peroxide methods. *J Med Plants Econ Dev*. 2020;4(1):a83. [[CrossRef](#)].
- Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep*. 2017;7:42717. [[CrossRef](#)].
- Vittaya L, Aiamyang S, Ui-eng J, Knongsai S, Leesakul N. Effect of solvent extraction on phytochemical component and antioxidant activity of vine and rhizome *Ampelocissus martini*. *Sci Technol Asia*. 2019;24(3):17–26. [[CrossRef](#)].
- Senguttuvan J, Paulsamy S, Karthika K. Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb, *Hypochaeris radicata* L. for *in vitro* antioxidant activities. *Asian Pac J Trop Biomed*. 2014;4:S359–67. [[CrossRef](#)].
- M02–A11. Performance standards for antimicrobial disk susceptibility tests. Wayne, PA, USA: Clinical and Laboratory Standards Institute; 2012.
- M100–S29. Performance standards for antimicrobial susceptibility testing. Wayne, PA, USA: Clinical and Laboratory Standards Institute; 2019.
- Campoccia D, Ravaoli S, Santi S, Mariani V, Santarcangelo C, De Filippis A, et al. Exploring the anticancer effects of standardized extracts of poplar-type Propolis: *in vitro* cytotoxicity toward cancer and normal cell lines. *Biomed Pharmacother*. 2021;141:111895. [[CrossRef](#)].
- Dai J, Mumper RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 2010;15(10):7313–52. [[CrossRef](#)].
- Zhang QW, Lin LG, Ye WC. Techniques for extraction and isolation of natural products: a comprehensive review. *Chin Med*. 2018;13:20. [[CrossRef](#)].

21. Henkel S, Misuraca MC, Troselj P, Davidson J, Hunter CA. Polarisation effects on the solvation properties of alcohols. *Chem Sci*. 2017;9(1):88–99. [[CrossRef](#)].
22. Nabeelah Bibi S, Fawzi MM, Gokhan Z, Rajesh J, Nadeem N, Rengasamy Kannan RR, et al. Ethnopharmacology, phytochemistry, and global distribution of Mangroves—a comprehensive review. *Mar Drugs*. 2019;17(4):231. [[CrossRef](#)].
23. Malik EM, Müller CE. Anthraquinones as pharmacological tools and drugs. *Med Res Rev*. 2016;36(4):705–48. [[CrossRef](#)].
24. Amala Dev AR, Sonia Mol J. Volatile chemical profiling and distinction of Citrus essential oils by GC analyses with correlation matrix; evaluation of its *in vitro* radical scavenging and microbicidal efficacy. *Results Chem*. 2024;7:101460. [[CrossRef](#)].
25. Ullah A, Munir S, Badshah SL, Khan N, Ghani L, Poulson BG, et al. Important flavonoids and their role as a therapeutic agent. *Molecules*. 2020;25(22):5243. [[CrossRef](#)].
26. Mlcek J, Jurikova T, Skrovankova S, Sochor J. Quercetin and its anti-allergic immune response. *Molecules*. 2016;21(5):623. [[CrossRef](#)].
27. Ganeshpurkar A, Saluja AK. The pharmacological potential of rutin. *Saudi Pharm J*. 2017;25(2):149–64. [[CrossRef](#)].
28. Vergara-Jimenez M, Almatrafi MM, Fernandez ML. Bioactive components in *Moringa oleifera* leaves protect against chronic disease. *Antioxidants*. 2017;6(4):91. [[CrossRef](#)].
29. Wang Z, Xu J, Wang Y, Xiang L, He X. Total saponins from *Tupistra chinensis* baker inhibits growth of human gastric cancer cells *in vitro* and *in vivo*. *J Ethnopharmacol*. 2021;278:114323. [[CrossRef](#)].
30. Kumar N, Goel N. Phenolic acids: natural versatile molecules with promising therapeutic applications. *Biotechnol Rep*. 2019;24:e00370. [[CrossRef](#)].
31. Divekar PA, Narayana S, Divekar BA, Kumar R, Gadratagi BG, Ray A, et al. Plant secondary metabolites as defense tools against herbivores for sustainable crop protection. *Int J Mol Sci*. 2022;23(5):2690. [[CrossRef](#)].
32. Lichtenthaler HK, Buschmann C. Chlorophylls and carotenoids: measurement and characterization by UV-VIS spectroscopy. *Curr Protoc Food Anal Chem*. 2001;1(1):F4.3.1–8. [[CrossRef](#)].
33. Schulz H, Baranska M. Identification and quantification of valuable plant substances by IR and Raman spectroscopy. *Vib Spectrosc*. 2007;43(1):13–25. [[CrossRef](#)].
34. Tella JO, Oseni SO. Comparative profiling of solvent-mediated phytochemical expressions in *Ocimum gratissimum* and *Vernonia amygdalina* leaf tissues via FTIR spectroscopy and colorimetric assays. *J Adv Med Pharm Sci*. 2019:1–25. [[CrossRef](#)].
35. Silverstein RM, Webster FX, Kiemle DJ. Spectrometric identification of organic compounds. 7th ed. Hoboken, NJ, USA: John Wiley & Sons, Inc.; 2005.
36. Smith BC. Infrared spectral interpretation: a systematic approach. 1st ed. Boca Raton, FL, USA: CRC Press; 1999. 288 p.
37. Bandaranayake WM. Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetl Ecol Manag*. 2002;10(6):421–52. [[CrossRef](#)].
38. Verma N, Shukla S. Impact of various factors responsible for fluctuation in plant secondary metabolites. *J Appl Res Med Aromat Plants*. 2015;2(4):105–13. [[CrossRef](#)].
39. Bento JAC, Ribeiro PRV, Bassinello PZ, de Brito ES, Zocollo GJ, Caliar M, et al. Phenolic and saponin profile in grains of carioca beans during storage. *LWT*. 2021;139:110599. [[CrossRef](#)].
40. Stalikas CD. Extraction, separation, and detection methods for phenolic acids and flavonoids. *J Sep Sci*. 2007;30(18):3268–95. [[CrossRef](#)].
41. Lavola A, Aphalo PJ, Lahti M, Julkunen-Tiitto R. Nutrient availability and the effect of increasing UV-B radiation on secondary plant compounds in Scots pine. *Environ Exp Bot*. 2003;49(1):49–60. [[CrossRef](#)].
42. Ramakrishna A, Ravishankar GA. Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal Behav*. 2011;6(11):1720–31. [[CrossRef](#)].
43. Dias MC, Pinto DCGA, Silva AMS. Plant flavonoids: chemical characteristics and biological activity. *Molecules*. 2021;26(17):5377. [[CrossRef](#)].
44. Kumari P, Gaur SS, Tiwari RK. Banana and its by-products: a comprehensive review on its nutritional composition and pharmacological benefits. *eFood*. 2023;4(5):e110. [[CrossRef](#)].
45. Leyva-Jimenez FJ, Lozano-Sanchez J, Borrás-Linares I, de la Luz Cadiz-Gurrea M, Mahmoodi-Khaledi E. Potential antimicrobial activity of honey phenolic compounds against Gram positive and Gram negative bacteria. *LWT*. 2019;101:236–45. [[CrossRef](#)].

46. Vadillo-Rodríguez V, Cavagnola MA, Pérez-Giraldo C, Fernández-Calderón MC. A physico-chemical study of the interaction of ethanolic extracts of Propolis with bacterial cells. *Colloids Surf B Biointerfaces*. 2021;200:111571. [[CrossRef](#)].
47. Goyal S, Thirumal D, Singh S, Kumar D, Singh I, Kumar G, et al. Basics of antioxidants and their importance. In: Sindhu RK, Singh I, Babu MA, editors. *Antioxidants: nature's defense against disease*. Hoboken, NJ, USA: John Wiley & Sons; 2025. p. 1–20. [[CrossRef](#)].
48. Rammohan A, Zyryanov GV, Bhagath YB, Manjula K. Antioxidants: structure-activity of plant polyphenolics. In: Litwack G, editor. *Vitamins and hormones*. San Diego, CA, USA: Academic Press; 2023. p. 395–411. [[CrossRef](#)].
49. Aribisala JO, Sabiu S. Redox impact on bacterial macromolecule: a promising avenue for discovery and development of novel antibacterials. *Biomolecules*. 2022;12(11):1545. [[CrossRef](#)].
50. Othman L, Sleiman A, Abdel-Massih RM. Antimicrobial activity of polyphenols and alkaloids in middle eastern plants. *Front Microbiol*. 2019;10:911. [[CrossRef](#)].
51. Maheshwari N, Sharma MC. Anticancer properties of some selected plant phenolic compounds: future leads for therapeutic development. *J Herb Med*. 2023;42:100801. [[CrossRef](#)].
52. Anand P, Sundaram C, Jhurani S, Kunnumakkara AB, Aggarwal BB. Curcumin and cancer: an “old-age” disease with an “age-old” solution. *Cancer Lett*. 2008;267(1):133–64. [[CrossRef](#)].
53. Bishayee A. Cancer prevention and treatment with resveratrol: from rodent studies to clinical trials. *Cancer Prev Res*. 2009;2(5):409–18. [[CrossRef](#)].
54. Bakun P, Mlynarczyk DT, Koczorowski T, Cerbin-Koczorowska M, Piwowarczyk L, Kolasiński E, et al. Tea-break with epigallocatechin gallate derivatives—powerful polyphenols of great potential for medicine. *Eur J Med Chem*. 2023;261:115820. [[CrossRef](#)].
55. Tabassum S, Rosli N, Ichwan SJA, Mishra P. Thymoquinone and its pharmacological perspective: a review. *Pharmacol Res Mod Chin Med*. 2021;1:100020. [[CrossRef](#)].
56. Ruangsuriya J, Sichaem J, Tantraworasin A, Saeteng S, Wongmaneerung P, Inta A, et al. Phytochemical profiles and anticancer effects of *Calophyllum inophyllum* L. extract relating to reactive oxygen species modulation on patient-derived cells from breast and lung cancers. *Scientifica*. 2023;2023:6613670. [[CrossRef](#)].
57. Mohammadi Nejad S, Özgüneş H, Başaran N. Pharmacological and toxicological properties of eugenol. *TJPS*. 2017;14(2):201–6. [[CrossRef](#)].
58. Ramírez-Santos J, Calzada F, García-Hernández N, Barbosa E, Velázquez C, Valdes M. Caryophyllene oxide, a bicyclic terpenoid isolated from *Annona macrophyllata* with antitumor activity: *in vivo*, *in vitro*, and *in silico* studies. *Int J Mol Sci*. 2024;25(24):13355. [[CrossRef](#)].
59. Sharma H, Sehgal R, Jhacak S, Deshmukh K, Nada R. Evaluation of farnesol orally and topically against experimental cutaneous leishmaniasis: *in-vivo* analysis. *PLoS One*. 2023;18(8):e0290297. [[CrossRef](#)].
60. Gori M, Altomare A, Cocca S, Solida E, Ribolsi M, Carotti S, et al. Palmitic acid affects intestinal epithelial barrier integrity and permeability *in vitro*. *Antioxidants*. 2020;9(5):417. [[CrossRef](#)].
61. Zhu J, Park S, Kim CH, Jeong KH, Kim WJ. Eugenol alleviates neuronal damage via inhibiting inflammatory process against pilocarpine-induced status epilepticus. *Exp Biol Med*. 2023;248(8):722–31. [[CrossRef](#)].
62. Fidyk K, Fiedorowicz A, Strzdała L, Szumny A. β -caryophyllene and β -caryophyllene oxide-natural compounds of anticancer and analgesic properties. *Cancer Med*. 2016;5(10):3007–17. [[CrossRef](#)].
63. Karakaya S, Yilmaz SV, Özdemir Ö, Koca M, Pinar NM, Demirci B, et al. A caryophyllene oxide and other potential anticholinesterase and anticancer agent in *Salvia verticillata* subsp. *amasiaca* (Freyn & Bornm.) Bornm. (Lamiaceae). *J Essent Oil Res*. 2020;32(6):512–25. [[CrossRef](#)].
64. Phonglo A, Dowerah D, Sarma S, Ahmed N, Dutta P, Basumutary M, et al. Essential oil constituents of regional ethnomedicinal plants as potential inhibitors of SARS-CoV-2 M(pro): an integrated molecular docking, molecular dynamics and QM/MM study. *J Biomol Struct Dyn*. 2024:1–23. [[CrossRef](#)].
65. Krishnaveni K, Murugan M, Kalaimathi R, Basha A, Pallan G, Kandeepan C, et al. ADMET informatics of Plant Derived n-Hexadecanoic Acid (Palmitic Acid) from ethyl acetate fraction of *Moringa oleifera* leaf extract. *J Drug Delivery Ther*. 2022;12(5):132–45. [[CrossRef](#)].
66. Zhang MQ, Wilkinson B. Drug discovery beyond the ‘rule-of-five’. *Curr Opin Biotechnol*. 2007;18(6):478–88. [[CrossRef](#)].