

Impact of SARS-CoV-2 infection during pregnancy on postnatal brain development: The potential role of glial cells

LARISSA DANIELE BOBERMIN^{1,*}; LARA SCOPEL MEDEIROS²; FERNANDA WEBER²; GIANCARLO TOMAZZONI DE OLIVEIRA²; LUCÉLIA SANTI^{3,4,5}; WALTER ORLANDO BEYS-DA-SILVA^{3,4,5}; CARLOS-ALBERTO GONÇALVES^{1,2,6}; ANDRÉ QUINCOZES-SANTOS^{1,2,6}

¹ Programa de Pós-Graduação em Neurociências, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, 90050170, Brazil

² Departamento de Bioquímica, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, 90035003, Brazil

³ Faculdade de Farmácia, Universidade Federal do Rio Grande do Sul, Porto Alegre, 90610000, Brazil

⁴ Hospital de Clínicas de Porto Alegre, Porto Alegre, 90035903, Brazil

⁵ Programa de Pós-Graduação em Biologia Celular e Molecular, Centro de Biotecnologia, Universidade Federal do Rio Grande do Sul, Porto Alegre, 91501970, Brazil

⁶ Programa de Pós-Graduação em Ciências Biológicas: Bioquímica, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, 90035003, Brazil

Key words: Brain development, Glial cells, Inflammation, Pregnancy, SARS-CoV-2

Abstract: Glial cells are crucial for maintaining central nervous system (CNS) homeostasis. They actively participate in immune responses, as well as form functional barriers, such as blood-brain barrier (BBB), which restrict the entry of pathogens and inflammatory mediators into the CNS. In general, viral infections during the gestational period can alter the embryonic and fetal environment, and the related inflammatory response may affect neurodevelopment and lead to behavioral dysfunction during later stage of life, as highlighted by our group for Zika virus infection. Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) induces a cytokine storm and, during pregnancy, may be related to a more severe form of the coronavirus disease-19 (COVID-19) and also to higher preterm birth rates. SARS-CoV-2 can also affect the CNS by inducing neurochemical remodeling in neural cells, which can compromise neuronal plasticity and synaptic function. However, the impact of SARS-CoV-2 infection during pregnancy on postnatal CNS, including brain development during childhood and adulthood, remains undetermined. Our group has recently highlighted the impact of COVID-19 on the expression of molecular markers associated with neuropsychiatric disorders, which are strongly related to the inflammatory response. Thus, based on these relationships, we discussed the impact of SARS-CoV-2 infection either during pregnancy or in critical periods of neurodevelopment as a risk factor for neurological consequences in the offspring later in life, focusing on the potential role of glial cells. Thus, it is important to consider future and long-term public health concerns associated with SARS-CoV-2 infection during pregnancy.

Introduction

New public health concerns have been raised by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection in pregnant women, which may be related to the development of a more severe form of the coronavirus disease-19 (COVID-19), gestational complications, and higher preterm birth rates (Beys-da-Silva *et al.*, 2020; Celewicz *et al.*, 2021; Karasek *et al.*, 2021). However, the impact of COVID-19 during pregnancy on postnatal development, particularly development of the

central nervous system (CNS), including during childhood and adulthood, remains undetermined. In general, viral infections during the gestational period can alter the embryonic and fetal environment, and the related inflammatory response may affect neurodevelopment and lead to subsequent behavioral dysfunctions (Zimmer *et al.*, 2021). While relevant evidence has suggested vertical transmission of SARS-CoV-2 infection, pro-inflammatory mediators involved in the maternal cytokine storm may also pass through the placenta, resulting in an indirect inflammatory response in the fetus (Joma *et al.*, 2021; Vivanti *et al.*, 2020).

Neurological complications have also been increasingly reported in individuals with COVID-19, which may be due to either direct viral entry into the CNS or a systemic cytokine storm (Doyle, 2021; Solomon, 2021). Angiotensin-converting

*Address correspondence to: Larissa Daniele Bobermin, larissa.bobermin@ufrgs.br

Received: 21 January 2022; Accepted: 18 April 2022



enzyme 2 (ACE-2) is expressed in glial cells and neurons, which supports the potential neurotropism of SARS-CoV-2 (Hernández *et al.*, 2021). Moreover, glial cells, particularly microglia and astrocytes, perform immune and inflammatory functions that allow them to respond to both viral infection and systemic inflammation (Liddelow *et al.*, 2020; Tremblay *et al.*, 2020). Importantly, glial cells also play crucial roles in neurodevelopment. In this context, the dysfunction of these cells during the fetal and postnatal periods may be related to neuropsychiatric disorders such as autism spectrum disorder (ASD) and schizophrenia, later in life (Dietz *et al.*, 2020; Scuderi and Verkhatsky, 2020). In addition, TORCH (an acronym used to refer to toxoplasmosis, rubella, cytomegalovirus, herpes simplex) congenital syndromes have been associated with several neurological complications (Megli and Coyne, 2021). In this article, we highlight the potential consequences of SARS-CoV-2 infection during pregnancy on brain development.

Immune, homeostatic, and neurodevelopmental roles of glial cells

Glial cells can be subdivided into macroglia (astrocytes and oligodendrocytes) and microglia, which are essential for brain development and homeostasis (Zuchero and Barres, 2015). Although microglial cells are resident immune cells in the CNS, astrocytes actively participate in innate immunity and inflammatory responses (Colombo and Farina, 2016). Both microglia and astrocytes synthesize and release several cytokines and chemokines that can propagate inflammation within the CNS (Liddelow *et al.*, 2020). Astrocytes possess a specific cytoarchitecture that allows intimate contact with blood vessels, synapses, and other glial cells, which enables them to sense their surroundings and dynamically respond to changes in their microenvironment. They form functional barriers, such as the blood-brain barrier (BBB), which restricts the entry of pathogens and inflammatory mediators into the CNS (Greenhalgh *et al.*, 2020). It is noteworthy that peripheral immune cells can affect the CNS when BBB permeability is compromised, and that astrocytes can guide inflammatory processes under healthy as well as diseased and injured conditions. Moreover, astrocytes regulate neurotransmitter systems, metabolic activity, and synthesis and release neurotrophic factors (Quincozes-Santos *et al.*, 2021b).

Macroglia are derived from radial glial cells that have a neuroepithelial origin and are pluripotent neural cell precursors (Zuchero and Barres, 2015). Microglial cells have a distinct (mesodermal) origin from the embryonic yolk sac and infiltrate the developing neural tube (Ginhoux and Prinz, 2015). The proliferation of glial and neuronal cells peaks during fetal development, between the 9th and 16th week post-conception. This means that glial cells are present during most brain development processes, including neurogenesis, neuronal circuit building during prenatal periods, and synaptic pruning in postnatal stages (Ziats *et al.*, 2015). While glial cells develop, proliferate, differentiate, and acquire functional properties, they establish important crosstalk with other glial cell types and neurons, which is continuously required not only during development but also in the mature brain. This communication occurs mainly via

secreted molecules, including trophic and growth factors, cytokines, chemokines, neurotransmitters, and hormones (Han *et al.*, 2021; Tay *et al.*, 2017).

During development, astrocytes are essential for synaptogenesis, production and delivery of glutamate to neurons, extracellular potassium buffering, and metabolic support, and can respond to neuronal activity. Thus, they display a wide range of homeostatic functions that are necessary for synapse formation and maintenance (Zuchero and Barres, 2015). Microglia are necessary for neurogenesis and neuronal migration, orchestrating the assembly of prenatal neuronal circuits (Colonna and Butovsky, 2017). During postnatal development, both microglia and astrocytes are responsible for tagging and eliminating redundant neurons that do not form functional neuronal circuits, a process known as synaptic pruning (Colonna and Butovsky, 2017).

Glial cells are at the center of CNS homeostasis and respond to damage. Disturbances in glial morphology, gene expression, and functions have been reported in a growing number of neurodevelopmental and neuropsychiatric disorders, such as ASD and schizophrenia (Bristol Silvestrin *et al.*, 2013; Dietz *et al.*, 2020; Gottfried *et al.*, 2015). While the precise time of onset and cellular neurobiology of these disorders remain poorly understood, the role of glial cells in these disorders has gained evidence. Different mechanisms have been proposed to link aberrant glial development/activity during gestational and postnatal periods to ASD and schizophrenia. Neuroinflammation is guided by morphological and functional changes in microglia and astrocytes. In this context, in utero exposure to pathogens and other neurotoxic environmental factors that cause neuroinflammation in the developing brain has emerged as an etiological factor for these disorders (Gottfried *et al.*, 2015; Scuderi and Verkhatsky, 2020). Moreover, non-inflammatory mechanisms may involve defects in synaptogenesis or synaptic pruning, disruption of glutamate metabolism, potassium homeostasis, secretion of neurotrophic factors, and abnormal glial coverage and synaptic support (Ballas *et al.*, 2009; Dietz *et al.*, 2020; Maezawa and Jin, 2010). All of these glial cell-associated failures may result in dysregulated synaptic transmission and/or a lack of functional neurons.

Maternal infections and later neurological consequences in the offspring: potential roles of glial cells

An important connection between maternal infections during pregnancy and congenital syndromes and/or later neuropsychiatric disorders has been established using epidemiological and experimental animal models (Estes and McAllister, 2016; Knuesel *et al.*, 2014). Gestational transmission can occur during the prenatal period through the transplacental hematogenous route, during the perinatal period through contact with blood and vaginal secretions from the mother, and during the postnatal period through breast milk (Neu *et al.*, 2015). Importantly, although evidence of infection can be observed at birth and/or in childhood, it is possible that certain symptoms might manifest only in adulthood.

TORCH refers to congenital syndromes with clinical similarities, namely infections caused by *Toxoplasma gondii*,

rubella, cytomegalovirus (CMV), and herpes simplex (HSV) (Megli and Coyne, 2021). The common clinical manifestations of TORCH congenital syndromes include microcephaly, cerebral calcifications, hydrocephalus, motor and brain dysfunction, sensorineural hearing loss, and intellectual disability (Megli and Coyne, 2021). CMV infection has been shown to alter the proliferation, migration, and differentiation of neural precursor cells, including astrocytes, which may explain the abnormalities in brain development observed in congenital infections (Odeberg *et al.*, 2007). Amendments in the term TORCH, such as TORCHES or STORCH, have been proposed to include syphilis, which is caused by *Treponema pallidum* (Kinney and Kumar, 1988). More recently, due to the severity of congenital infection, Zika virus (ZIKV) infection has also been proposed to be included as a new member, as TORCHZ or STORCHZ (Coyne and Lazear, 2016).

Although ZIKV infection during pregnancy is mainly associated with microcephaly, it has recently been correlated with ASD diagnosis in childhood (Santi *et al.*, 2021; Vianna *et al.*, 2018). These neurodevelopmental abnormalities may be associated with neuroimmune changes caused by the release of inflammatory mediators by ZIKV-infected neural cells (Bobermin *et al.*, 2020). Moreover, fetal human astrocytes can be used as a viral reservoir, contributing to chronic brain infections (Limonta *et al.*, 2018). Other viruses, such as rubella virus and influenza, can also target glial cells and interfere with neuron-glia communication, resulting in aberrant neuronal migration, synaptogenesis, and synaptic pruning associated with ASD-like behaviors and an increased risk for schizophrenia (Fatemi *et al.*, 2008; Scuderi and Verkhatsky, 2020). It is important to note that several neuropsychiatric disorders, other than STORCHZ, have also been reported to be associated with morphofunctional changes.

Of note, both direct infection by viruses in the fetal CNS and maternal immune responses may be critical for these neurodevelopmental and/or neuropsychiatric sequelae. Activation of Toll-like receptor (TLR) pathways (including TLR3) in host immune cells, which may include glial cells, induces the expression of inflammatory cytokines and interferons through nuclear factor κ B (NF κ B) signaling (Alexopoulou *et al.*, 2001; Greenhalgh *et al.*, 2020). In this context, pro-inflammatory cytokines such as interleukins (IL-6, IL-1 β , IL-17) and tumor necrosis factor α (TNF- α) have been described in animal models as key signaling molecules that predispose the individual to neurodevelopmental and neuropsychiatric disorders (Favrais *et al.*, 2011; Gilmore *et al.*, 2005).

Current evidence on SARS-CoV-2 vertical transmission

Relevant evidence has raised the possibility of SARS-CoV-2 vertical transmission, either during the antepartum, intrapartum, or postpartum periods. ACE-2 expression in the placenta and detection of SARS-CoV-2 in the placenta and amniotic fluid suggest that transplacental transmission can occur (Hosier *et al.*, 2020; Li *et al.*, 2020; Vivanti *et al.*, 2020). Pathological findings have also shown increased inflammatory infiltration in the placentas of SARS-CoV-2-infected pregnant women (Schoenmakers *et al.*, 2021). Notably, the placenta may become more susceptible to SARS-CoV-2 infection when it is inflamed (Lye *et al.*, 2020).

Moreover, SARS-CoV-2 has been detected in the umbilical cord, vaginal mucosa, and milk (Fenizia *et al.*, 2020).

In neonates born from SARS-CoV-2-infected women, severe respiratory symptoms are not common, although there are some cases of respiratory diseases, abnormal Apgar indexes, and pneumonia (Nayak *et al.*, 2020; Wu *et al.*, 2020). However, a case report documented viremia and neurological symptoms in a newborn, which, in addition to supporting the transplacental transmission of SARS-CoV-2, suggested the effects of SARS-CoV-2 infection in the immature brain with evidence of gliosis (Vivanti *et al.*, 2020).

Neurological symptoms associated with SARS-CoV-2 infection: Potential role of glial cells and possible implications in neurodevelopment

Neurological manifestations are increasingly common in COVID-19 adult patients in addition to some degree of encephalitis, which could indicate substantial damage to the CNS (Doyle, 2021; Moriguchi *et al.*, 2020). Moreover, the development of neurological symptoms in children with COVID-19 has been described (Abdel-Mannan *et al.*, 2020). Although neurotropism and the mechanisms associated with SARS-CoV-2 entry into the CNS are still under investigation, it is important to note that other coronaviruses can compromise the nervous system by directly targeting neural cells and inducing a significant inflammatory condition that can lead to neuronal death (Li *et al.*, 2016).

In line with this, it has been demonstrated that the main cells of the CNS, such as neurons, astrocytes, and microglia, may be direct targets of SARS-CoV-2 owing to the expression of ACE-2 (Cama *et al.*, 2021; Crunfli *et al.*, 2020; Song *et al.*, 2021). In particular, astrocytes have been shown to be permissive to SARS-CoV-2, serving as a potential site for replication of the virus in the CNS. Moreover, astrocytes undergo several functional alterations after infection, including metabolic remodeling, which can compromise neuronal plasticity and synaptic functions (Crunfli *et al.*, 2020). Therefore, astrocyte dysfunction may be a mechanism underlying the short- and long-term neurological consequences of SARS-CoV-2 infection.

Reactive astrogliosis has also been observed in post-mortem brains of COVID-19 patients, along with microglial activation and infiltration of T lymphocytes, reinforcing the presence of an inflammatory process (Matschke *et al.*, 2020). Notably, both astrocytes and microglia are sensitive to peripheral cytokine storms, which can disrupt the BBB and facilitate viral entry into the CNS (Greenhalgh *et al.*, 2020; Huang *et al.*, 2021). Importantly, glial inflammatory phenotypes may contribute to various brain dysfunctions, including neurodegenerative diseases, neuropsychiatric symptoms, and neurodevelopmental disorders (Garden and Campbell, 2016; Pantelis *et al.*, 2020).

The cytokine storm caused by SARS-CoV-2 may be an aggravating factor in pregnant women, since maternal pro-inflammatory mediators can be transferred across the placenta and subsequently reach the developing CNS (Joma *et al.*, 2021; Reyes-Lagos *et al.*, 2021). In this regard, high levels of important inflammatory markers, including IL-6, IL-17, and TNF- α , have been reported in the blood of pregnant women with COVID-19 and the umbilical cord, as

well as in the serum of neonates born to infected mothers (Joma *et al.*, 2021; Xiao *et al.*, 2020; Zhu *et al.*, 2020). Among them, the role of IL-6 in maternal systemic inflammation during fetal brain development has been particularly studied. In an animal model, maternal treatment with IL-6 alone was sufficient to predispose the offspring to neuropsychiatric disorders such as schizophrenia and autism (Smith *et al.*, 2007). Interestingly, inhibition of IL-6 signaling through its placental receptors prevented increased IL-6 expression in the fetal brain and neurodevelopmental abnormalities (Wu *et al.*, 2017). Moreover, studies in humans have shown architectural and functional brain alterations in children exposed to increased levels of IL-6 during pregnancy (Rasmussen *et al.*, 2019; Rudolph *et al.*, 2018). Therefore, since IL-6 is an important component of the cytokine storm in COVID-19, it might act as a crucial molecular mediator in the potential link between SARS-CoV-2 infection during pregnancy and neurodevelopmental and neuropsychiatric disorders in the offspring.

It is also important to consider that TNF- α and IL-17, in addition to other inflammatory mediators whose levels are commonly increased in COVID-19 patients (C-reactive protein, IL-1 β , IL-2, IL-8, and interferons), may contribute to neurodevelopmental and neuropsychiatric diseases related to maternal immune activation (Jiang *et al.*, 2018). Although the mechanisms by which cytokines affect brain development remain poorly understood, they seem to involve changes in functional brain connectivity, white matter damage, and abnormalities in neurotransmission (Jiang *et al.*, 2018; Mirabella *et al.*, 2021; Saunders *et al.*, 2020). Cytokines from the periphery can also induce CNS responses in several ways, including diffusion across the disrupted BBB and induction of synthesis and release of cytokines by glial cells (Gu *et al.*, 2021; Jiang *et al.*, 2018). Therefore, even in the absence of vertical transmission or direct neuroinvasion by SARS-CoV-2, changes in the fetal and neonatal brain functions may occur due to immune activation.

In addition to environmental factors, susceptibility to neurodevelopmental and neuropsychiatric disorders may also involve molecular factors. Defects in glial genes related to synaptogenesis, synaptic transmission, and neuroinflammation have been reported in patients diagnosed with ASD (Scuderi and Verkhatsky, 2020). Supporting the role of altered glial-neuronal relationships during fetal CNS development in ASD, some genes that are essential for brain development and highly expressed in different glial cell types were also found to confer susceptibility to ASD (Scuderi and Verkhatsky, 2020). Notably, an analysis of differential gene expression datasets for clinical samples of COVID-19 patients identified a high number of genes that are associated with the pathophysiology of several neuropsychiatric disorders, including ASD, schizophrenia, bipolar disorder, depression, and Alzheimer's disease (Quincozes-Santos *et al.*, 2021a; Quincozes-Santos *et al.*, 2021).

Concluding Remarks

Glial cells are important players in neurodevelopment and maintenance of brain homeostasis, and disruption of their




FIGURE 1. Putative role of glial cells in the possible neurodevelopmental and/or neuropsychiatric consequences related to SARS-CoV-2 infection during pregnancy. Recent evidence has raised the possibility of vertical transmission of SARS-CoV-2 as well as a direct brain infection. Additionally, maternal cytokine storm related to COVID-19 may cause fetal neuroinflammation. Both a direct infection and an inflammatory response triggered by SARS-CoV-2 might impact brain development by affecting critical functions of glial cells, as it has been demonstrated for other viral infections. These altered glial functions may increase the risk of later neurodevelopmental/neuropsychiatric disorders. The cells in the circle represent functional glial cells and neurons. Astrocytes are represented in blue; microglia are represented in yellow; SARS-CoV-2 is represented in green and cytokines/immune cells in pink.

functions has been strongly associated with a wide range of neuropsychiatric disorders. The relationship between maternal infections and possible neurodevelopmental disorders and/or later neuropsychiatric consequences in the offspring has been established, in which the available evidence supports glial-related pathogenesis. Based on these relationships, SARS-CoV-2 infection either during pregnancy or during critical neurodevelopmental periods could be related to an increased risk of neurological consequences in offsprings later in life (Fig. 1); thus, efforts to unravel this hypothesis are currently needed. It is also possible to conceive expansion of the current TORCH/STORCHZ family, generating a new acronym to incorporate other emerging pathogens. Moreover, it is also important to consider the future and long-term public health concerns associated with SARS-CoV-2 infection during pregnancy, as well as a careful follow-up of infants born to COVID-19 positive mothers.

Author Contribution: The authors confirm contribution to the paper as follows: LDB, LSM, FW, GTO, LS, WOBS, CAG, and AQS are responsible for conception, design, and draft manuscript preparation. All authors reviewed and approved the final version of the manuscript.

Ethics Approval: Not applicable.

Funding Statement: The authors are supported by Universidade Federal do Rio Grande do Sul (UFRGS), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS), and Instituto Nacional de Ciência e Tecnologia para Excitotoxicidade e Neuroproteção (INCTEN/CNPq).

Conflicts of Interest: The authors declare that they have no conflicts of interest to report regarding the present study.

References

- Abdel-Mannan O, Eyre M, Löbel U, Bamford A, Eltz C, Hameed B, Hemingway C, Hachohen Y (2020). Neurologic and radiographic findings associated with COVID-19 infection in children. *JAMA Neurology* **77**: 1440. DOI 10.1001/jamaneurol.2020.2687.
- Alexopoulou L, Holt AC, Medzhitov R, Flavell RA (2001). Recognition of double-stranded RNA and activation of NF- κ B by Toll-like receptor 3. *Nature* **413**: 732–738. DOI 10.1038/35099560.
- Ballas N, Liou DT, Grunseich C, Mandel G (2009). Non-cell autonomous influence of MeCP2-deficient glia on neuronal dendritic morphology. *Nature Neuroscience* **12**: 311–317. DOI 10.1038/nn.2275.
- Beys-da-Silva WO, da Rosa RL, Santi L, Tureta EF, Terraciano PB et al. (2020). The risk of COVID-19 for pregnant women: Evidences of molecular alterations associated with preeclampsia in SARS-CoV-2 infection. *Biochimica et Biophysica Acta (BBA) Molecular Basis of Disease* **1867**: 165999. DOI 10.1016/j.bbadis.2020.165999.
- Bobermin LD, Quincozes-Santos A, Santos CL, Varela APM, Teixeira TF et al. (2020). Zika virus exposure affects neuron-glia communication in the hippocampal slices of adult rats. *Scientific Reports* **10**: 21604. DOI 10.1038/s41598-020-78735-y.
- Bristol Silvestrin R, Bambini-Junior V, Galland F, Daniele Bobermin L, Quincozes-Santos A et al. (2013). Animal model of autism induced by prenatal exposure to valproate: Altered glutamate metabolism in the hippocampus. *Brain Research* **1495**: 52–60. DOI 10.1016/j.brainres.2012.11.048.
- Cama VF, Marín-Prida J, Acosta-Rivero N, Acosta EF, Díaz LO et al. (2021). The microglial NLRP3 inflammasome is involved in human SARS-CoV-2 cerebral pathogenicity: A report of three post-mortem cases. *Journal of Neuroimmunology* **361**: 577728. DOI 10.1016/j.jneuroim.2021.577728.
- Celewicz A, Celewicz M, Michalczyk M, Woźniakowska-Gondek P, Krejczyk K, Misiak M, Rzepka R (2021). Pregnancy as a risk factor of severe COVID-19. *Journal of Clinical Medicine* **10**: 5458. DOI 10.3390/jcm10225458.
- Colombo E, Farina C (2016). Astrocytes: Key regulators of neuroinflammation. *Trends Immunology* **37**: 608–620. DOI 10.1016/j.it.2016.06.006.
- Colonna M, Butovsky O (2017). Microglia function in the central nervous system during health and neurodegeneration. *Annual Review of Immunology* **35**: 441–468. DOI 10.1146/annurev-immunol-051116-052358.
- Coyne CB, Lazear HM (2016). Zika virus—reigniting the TORCH. *Nature Reviews Microbiology* **14**: 707–715. DOI 10.1038/nrmicro.2016.125.
- Crunfli F, Carregari VC, Veras FP, Vendramini PH, Valença AGF et al. (2020). SARS-CoV-2 infects brain astrocytes of COVID-19 patients and impairs neuronal viability (preprint). *Neurology*. DOI 10.1101/2020.10.09.20207464.
- Dietz AG, Goldman SA, Nedergaard M (2020). Glial cells in schizophrenia: A unified hypothesis. *The Lancet Psychiatry* **7**: 272–281. DOI 10.1016/S2215-0366(19)30302-5.
- Doyle MF (2021). Central nervous system outcomes of COVID-19. *Translational Research* **241**: 41–51. DOI 10.1016/j.trsl.2021.09.002.
- Estes ML, McAllister AK (2016). Maternal immune activation: Implications for neuropsychiatric disorders. *Science* **353**: 772–777. DOI 10.1126/science.aag3194.
- Fatemi SH, Folsom TD, Reutiman TJ, Sidwell RW (2008). Viral regulation of aquaporin 4, connexin 43, microcephalin and nucleolin. *Schizophrenia Research* **98**: 163–177. DOI 10.1016/j.schres.2007.09.031.
- Favrais G, van de Looij Y, Fleiss B, Ramanantsoa N, Bonnin P et al. (2011). Systemic inflammation disrupts the developmental program of white matter. *Annals of Neurology* **70**: 550–565. DOI 10.1002/ana.22489.
- Fenzia C, Biasin M, Cetin I, Vergani P, Mileto D et al. (2020). Analysis of SARS-CoV-2 vertical transmission during pregnancy. *Nature Communications* **11**: 5128. DOI 10.1038/s41467-020-18933-4.
- Garden GA, Campbell BM (2016). Glial biomarkers in human central nervous system disease: Glial biomarkers in human CNS disease. *Glia* **64**: 1755–1771. DOI 10.1002/glia.22998.
- Gilmore JH, Jarskog LF, Vadlamudi S (2005). Maternal poly I: C exposure during pregnancy regulates TNF α , BDNF, and NGF expression in neonatal brain and the maternal-fetal unit of the rat. *Journal of Neuroimmunology* **159**: 106–112. DOI 10.1016/j.jneuroim.2004.10.008.
- Ginhoux F, Prinz M (2015). Origin of microglia: Current concepts and past controversies. *Cold Spring Harbor Perspectives in Biology* **7**: a020537. DOI 10.1101/cshperspect.a020537.
- Gottfried C, Bambini-Junior V, Francis F, Riesgo R, Savino W (2015). The impact of neuroimmune alterations in autism spectrum disorder. *Frontiers in Psychiatry* **6**: 121. DOI 10.3389/fpsy.2015.00121.
- Greenhalgh AD, David S, Bennett FC (2020). Immune cell regulation of glia during CNS injury and disease. *Nature Reviews Neuroscience* **21**: 139–152. DOI 10.1038/s41583-020-0263-9.
- Gu M, Mei XL, Zhao YN (2021). Sepsis and cerebral dysfunction: BBB damage, neuroinflammation, oxidative stress, apoptosis and autophagy as key mediators and the potential therapeutic approaches. *Neurotoxicity Research* **39**: 489–503. DOI 10.1007/s12640-020-00270-5.
- Han RT, Kim RD, Molofsky AV, Liddel SA (2021). Astrocyte-immune cell interactions in physiology and pathology. *Immunity* **54**: 211–224. DOI 10.1016/j.immuni.2021.01.013.
- Hernández VS, Zetter MA, Guerra EC, Hernández-Araiza I, Karuzin N, Hernández-Pérez OR, Eiden LE, Zhang L (2021). ACE2 expression in rat brain: Implications for COVID-19 associated neurological manifestations. *Experimental Neurology* **345**: 113837. DOI 10.1016/j.expneurol.2021.113837.
- Hosier H, Farhadian SF, Morotti RA, Deshmukh U, Lu-Culligan A et al. (2020). SARS-CoV-2 infection of the placenta. *Journal of Clinical Investigation* **130**: 4947–4953. DOI 10.1172/JCI139569.
- Huang X, Hussain B, Chang J (2021). Peripheral inflammation and blood-brain barrier disruption: Effects and mechanisms. *CNS Neuroscience & Therapeutics* **27**: 36–47. DOI 10.1111/cns.13569.

- Jiang NM, Cowan M, Moonah SN, Petri WA (2018). The impact of systemic inflammation on neurodevelopment. *Trends in Molecular Medicine* **24**: 794–804. DOI 10.1016/j.molmed.2018.06.008.
- Joma M, Fovet CM, Seddiki N, Gressens P, Laforge M (2021). COVID-19 and pregnancy: Vertical transmission and inflammation impact on newborns. *Vaccines 95-Molecular Approaches to the Control of Infectious Diseases* **9**: 391. DOI 10.3390/vaccines9040391.
- Karasek D, Baer RJ, McLemore MR, Bell AJ, Blebu BE et al. (2021). The association of COVID-19 infection in pregnancy with preterm birth: A retrospective cohort study in California. *The Lancet Regional Health–Americas* **2**: 100027. DOI 10.1016/j.lana.2021.100027.
- Kinney JS, Kumar ML (1988). Should we expand the TORCH complex? A description of clinical and diagnostic aspects of selected old and new agents. *Clinics in Perinatology* **15**: 727–744. DOI 10.1016/S0095-5108(18)30670-5.
- Knuesel I, Chicha L, Britschgi M, Schobel SA, Bodmer M, Hellings JA, Toovey S, Prinssen EP (2014). Maternal immune activation and abnormal brain development across CNS disorders. *Nature Reviews Neurology* **10**: 643–660. DOI 10.1038/nrneurol.2014.187.
- Li K, Wohlford-Lenane C, Perlman S, Zhao J, Jewell AK, Reznikov LR, Gibson-Corley KN, Meyerholz DK, McCray PB (2016). Middle east respiratory syndrome coronavirus causes multiple organ damage and lethal disease in mice transgenic for human dipeptidyl peptidase 4. *The Journal of Infectious Diseases* **213**: 712–722. DOI 10.1093/infdis/jiv499.
- Li M, Chen L, Zhang J, Xiong C, Li X (2020). The SARS-CoV-2 receptor ACE2 expression of maternal-fetal interface and fetal organs by single-cell transcriptome study. *PLoS One* **15**: e0230295. DOI 10.1371/journal.pone.0230295.
- Liddel SA, Marsh SE, Stevens B (2020). Microglia and astrocytes in disease: Dynamic duo or partners in crime? *Trends in Immunology* **41**: 820–835. DOI 10.1016/j.it.2020.07.006.
- Limonta D, Jovel J, Kumar A, Airo A, Hou S et al. (2018). Human fetal astrocytes infected with zika virus exhibit delayed apoptosis and resistance to interferon: Implications for persistence. *Viruses* **10**: 646. DOI 10.3390/v10110646.
- Lye P, Dunk CE, Zhang J, Wei Y, Nakpu J et al. (2020). SARS-CoV-2 cell entry gene ACE2 expression in immune cells that infiltrate the placenta in infection-associated preterm birth (preprint). *Obstetrics and Gynecology*. DOI 10.1101/2020.09.27.20201590.
- Maezawa I, Jin LW (2010). Rett syndrome microglia damage dendrites and synapses by the elevated release of glutamate. *Journal of Neuroscience* **30**: 5346–5356. DOI 10.1523/JNEUROSCI.5966-09.2010.
- Matschke J, Lütgehetmann M, Hagel C, Sperhake JP, Schröder AS et al. (2020). Neuropathology of patients with COVID-19 in Germany: A post-mortem case series. *The Lancet Neurology* **19**: 919–929. DOI 10.1016/S1474-4422(20)30308-2.
- Megli CJ, Coyne CB (2021). Infections at the maternal-fetal interface: An overview of pathogenesis and defence. *Nature Reviews Microbiology* **20**: 67–82. DOI 10.1038/s41579-021-00610-y.
- Mirabella F, Desiato G, Mancinelli S, Fossati G, Rasile M et al. (2021). Prenatal interleukin 6 elevation increases glutamatergic synapse density and disrupts hippocampal connectivity in offspring. *Immunity* **54**: 2611–2631.e8. DOI 10.1016/j.immuni.2021.10.006.
- Moriguchi T, Harii N, Goto J, Harada D, Sugawara H et al. (2020). A first case of meningitis/encephalitis associated with SARS-Coronavirus-2. *International Journal of Infectious Diseases* **94**: 55–58. DOI 10.1016/j.ijid.2020.03.062.
- Nayak AH, Kapote DS, Fonseca M, Chavan N, Mayekar R, Sarmalkar M, Bawa A (2020). Impact of the Coronavirus infection in pregnancy: A preliminary study of 141 patients. *The Journal of Obstetrics and Gynecology of India* **70**: 256–261. DOI 10.1007/s13224-020-01335-3.
- Neu N, Duchon J, Zachariah P (2015). TORCH infections. *Clinics in Perinatology* **42**: 77–103. DOI 10.1016/j.clp.2014.11.001.
- Odeberg J, Wolmer N, Falci S, Westgren M, Sundström E, Seiger Å., Söderberg-Nauclér C (2007). Late human cytomegalovirus (HCMV) proteins inhibit differentiation of human neural precursor cells into astrocytes. *Journal of Neuroscience Research* **85**: 583–593. DOI 10.1002/(ISSN)1097-4547.
- Pantelis C, Jayaram M, Hannan AJ, Wesselingh R, Nithianantharajah J et al. (2020). Neurological, neuropsychiatric and neurodevelopmental complications of COVID-19. *Australian & New Zealand Journal of Psychiatry* **55**: 750–762. DOI 10.1177/0004867420961472.
- Quincozes-Santos A, da Rosa RL, Bobermin LD, Tureta EF, Santi L et al. (2021). Association between molecular markers of COVID-19 and Alzheimer's disease. *Journal of Medical Virology*, DOI 10.1002/jmv.27391.
- Quincozes-Santos A, Rosa RL, Tureta EF, Bobermin LD, Berger M, Guimarães JA, Santi L, Beys-da-Silva WO (2021a). COVID-19 impacts the expression of molecular markers associated with neuropsychiatric disorders. *Brain, Behavior, & Immunity-Health* **11**: 100196. DOI 10.1016/j.bbih.2020.100196.
- Quincozes-Santos A, Santos CL, de Souza Almeida RR, da Silva A, Thomaz NK et al. (2021b). Gliotoxicity and glioprotection: The dual role of glial cells. *Molecular Neurobiology* **58**: 6577–6592. DOI 10.1007/s12035-021-02574-9.
- Rasmussen JM, Graham AM, Entringer S, Gilmore JH, Styner M, Fair DA, Wadhwa PD, Buss C (2019). Maternal interleukin-6 concentration during pregnancy is associated with variation in frontolimbic white matter and cognitive development in early life. *NeuroImage* **185**: 825–835. DOI 10.1016/j.neuroimage.2018.04.020.
- Reyes-Lagos JJ, Abarca-Castro EA, Echeverría JC, Mendieta-Zerón H, Vargas-Caraveo A, Pacheco-López G (2021). A translational perspective of maternal immune activation by SARS-CoV-2 on the potential prenatal origin of neurodevelopmental disorders: The role of the cholinergic anti-inflammatory pathway. *Frontiers in Psychology* **12**: 614451. DOI 10.3389/fpsyg.2021.614451.
- Rudolph MD, Graham AM, Feczko E, Miranda-Dominguez O, Rasmussen JM, Nardos R, Entringer S, Wadhwa PD, Buss C, Fair DA (2018). Maternal IL-6 during pregnancy can be estimated from newborn brain connectivity and predicts future working memory in offspring. *Nature Neuroscience* **21**: 765–772. DOI 10.1038/s41593-018-0128-y.
- Santi L, Riesgo RS, Quincozes-Santos A, Schuler-Faccini L, Tureta EF et al. (2021). Zika virus infection associated with autism spectrum disorder: A case report. *Neuroimmunomodulation* **28**: 229–232. DOI 10.1159/000516560.
- Saunders JM, Moreno JL, Ibi D, Sikaroodi M, Kang DJ et al. (2020). Gut microbiota manipulation during the prepubertal period shapes behavioral abnormalities in a mouse neurodevelopmental disorder model. *Scientific Reports* **10**: 4697. DOI 10.1038/s41598-020-61635-6.
- Schoenmakers S, Snijder P, Verdijk RM, Kuiken T, Kamphuis SSM et al. (2021). Severe acute respiratory syndrome Coronavirus 2 placental infection and inflammation leading to fetal distress and neonatal multi-organ failure in an asymptomatic woman.

- Journal of the Pediatric Infectious Diseases Society* **10**: 556–561. DOI 10.1093/jpids/piaa153.
- Scuderi C, Verkhratsky A (2020). The role of neuroglia in autism spectrum disorders. *Progress in Molecular Biology and Translational Science* **173**: 301–330. DOI 10.1016/bs.pmbts.2020.04.011.
- Smith SEP, Li J, Garbett K, Mirnics K, Patterson PH (2007). Maternal immune activation alters fetal brain development through interleukin-6. *Journal of Neuroscience* **27**: 10695–10702. DOI 10.1523/JNEUROSCI.2178-07.2007.
- Solomon T (2021). Neurological infection with SARS-CoV-2-the story so far. *Nature Reviews Neurology* **17**: 65–66. DOI 10.1038/s41582-020-00453-w.
- Song E, Zhang C, Israelow B, Lu-Culligan A, Prado AV et al. (2021). Neuroinvasion of SARS-CoV-2 in human and mouse brain. *Journal of Experimental Medicine* **218**: e20202135. DOI 10.1084/jem.20202135.
- Tay TL, Savage JC, Hui CW, Bisht K, Tremblay MÈ (2017). Microglia across the lifespan: From origin to function in brain development, plasticity and cognition: Microglia across the lifespan. *The Journal of Physiology* **595**: 1929–1945. DOI 10.1113/JP272134.
- Tremblay ME, Madore C, Bordeleau M, Tian L, Verkhratsky A (2020). Neuropathobiology of COVID-19: The role for glia. *Frontiers in Cellular Neuroscience* **14**: 592214. DOI 10.3389/fncel.2020.592214.
- Vianna P, do Gomes JA, Boquett JA, Fraga LR, Schuch JB, Vianna FSL, Schuler-Faccini L (2018). Zika virus as a possible risk factor for autism spectrum disorder: Neuroimmunological aspects. *Neuroimmunomodulation* **25**: 320–327. DOI 10.1159/000495660.
- Vivanti AJ, Vauloup-Fellous C, Prevot S, Zupan V, Suffee C, Do Cao J, Benachi A, de Luca D (2020). Transplacental transmission of SARS-CoV-2 infection. *Nature Communications* **11**: 3572. DOI 10.1038/s41467-020-17436-6.
- Wu WL, Hsiao EY, Yan Z, Mazmanian SK, Patterson PH (2017). The placental interleukin-6 signaling controls fetal brain development and behavior. *Brain, Behavior, and Immunity* **62**: 11–23. DOI 10.1016/j.bbi.2016.11.007.
- Wu YT, Liu J, Xu JJ, Chen YF, Yang W et al. (2020). Neonatal outcome in 29 pregnant women with COVID-19: A retrospective study in Wuhan, China. *PLoS Medicine* **17**: e1003195. DOI 10.1371/journal.pmed.1003195.
- Xiao T, Xia S, Zeng L, Lin G, Wei Q et al. (2020). A multicentre observational study on neonates exposed to SARS-CoV-2 in China: The Neo-SARS-CoV-2 study protocol. *BMJ Open* **10**: e038004. DOI 10.1136/bmjopen-2020-038004.
- Zhu H, Wang L, Fang C, Peng S, Zhang L, Chang G, Xia S, Zhou W (2020). Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Translational Pediatrics* **9**: 51–60. DOI 10.21037/tp.2020.02.06.
- Ziats MN, Edmonson C, Rennert OM (2015). The autistic brain in the context of normal neurodevelopment. *Frontiers in Neuroanatomy* **9**: 115. DOI 10.3389/fnana.2015.00115.
- Zimmer A, Youngblood A, Adnane A, Miller BJ, Goldsmith DR (2021). Prenatal exposure to viral infection and neuropsychiatric disorders in offspring: A review of the literature and recommendations for the COVID-19 pandemic. *Brain, Behavior, and Immunity* **91**: 756–770. DOI 10.1016/j.bbi.2020.10.024.
- Zuchero JB, Barres BA (2015). Glia in mammalian development and disease. *Development* **142**: 3805–3809. DOI 10.1242/dev.129304.