

Fatty acids in microalgae and cyanobacteria in a changing world: Contrasting temperate and cold environments

MARCELO P. HERNANDO^{1,2}; IRENE R. SCHLOSS^{3,4,5,*}; FLORENCIA DE LA ROSA^{6,7}; MARLEEN DE TROCH⁸

¹ Comisión Nacional de Energía Atómica (CNEA), Centro Atómico Constituyentes, Departamento de Radiobiología, San Martín, Buenos Aires, Argentina

² Red de Investigación de Estresores Marinos-costeros en América Latina y el Caribe (REMARCO), Mar del Plata, 7602, Argentina

³ Instituto Antártico Argentino (IAA), San Martín, Buenos Aires, Argentina

⁴ Centro Austral de Investigaciones Científicas (CADIC-CONICET), Ushuaia, 9410, Argentina

⁵ Universidad Nacional de Tierra del Fuego, Ushuaia, 9410, Argentina

⁶ Instituto de Ciencias Básicas y Experimentales (ICBE), Universidad de Morón, Morón, Buenos Aires, Argentina

⁷ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina

⁸ Ghent University, Faculty of Sciences, Biology Department, Marine Biology, Ghent, 9000, Belgium

Key words: Antarctic, Temperate, Essential FAs, Increased temperature, Microalgae, Cyanobacteria

Abstract: Under the present changing climate conditions and the observed temperature increase, it is of high importance to understand its effects on aquatic microbial life, and organisms' adaptations at the biochemical level. To adjust to temperature or salinity stress and avoid cell damage, organisms alter their degree of fatty acids (FAs) saturation. Thus, temperature is expected to have strong effects on both the quantity and quality of FAs in aquatic microorganisms. Here we review some recent findings about FAs sensitivity to climate change in contrasting environments. Overall, heat waves may induce changes in the relative abundance of polyunsaturated FAs (PUFA). However, the impact of the exposure to warming waters is different in temperate and polar environments. In cold marine waters, high concentration of omega-3 (ω 3) FAs such as eicosapentaenoic acid (EPA) is promoted due to the activation of the desaturase enzyme. In this way, cells have enough energy to produce or activate antioxidant protection mechanisms and avoid oxidative stress due to heat waves. Contrastingly, under high irradiance and heat wave conditions in temperate environments, photosystems' protection is achieved by decreasing EPA concentration due to desaturase sensitivity. Essential FAs are transferred in aquatic food webs. Therefore, any alteration in the production of essential FAs by phytoplankton (the main source of ω 3) due to climate warming can be transferred to higher trophic levels, with cascading effects for the entire aquatic ecosystem.

Introduction

Increasing fossil fuel emissions during the past decades have increased atmospheric CO₂ concentrations and lead to a rise of global average temperatures (Myhre *et al.*, 2015; Abhilash, 2015). The average surface temperature of the planet increased around 1°C during 2019, being the second warmest year in the 140-year record (NOAA, 2020). Temperature is expected to further increase globally with larger changes at higher latitudes (Perkins-Kirkpatrick and Gibson, 2017). Predictions made by the Intergovernmental Panel on Climate Change (IPCC, 2019) indicate that the surface temperature would increase approximately from 3 to

5°C by 2100. Furthermore, the average ocean temperature increased by 0.06°C per decade in the last 50 years (Lindsey and Dahlman, 2017). As a result of the increased surface and air temperatures and the frequency of extreme heat days, the marine and freshwater temperatures will also continue to rise (IPCC, 2019).

Aquatic ecosystems such as large shallow lakes are particularly vulnerable to climate change (van Doorslaer *et al.*, 2007). For example, it could produce substantial coastline changes (Pussella *et al.*, 2015). However, the increases in temperature vary regionally.

The physico-chemical properties of polar environments are shifting rapidly, especially in the Arctic and in areas around the Antarctic Peninsula. In the Arctic Ocean, a 2°C increase was observed over the last 20 years (Economist, 2013). Temperature records for the West Antarctic Peninsula

*Address correspondence to: Irene R. Schloss, ireschloss@gmail.com
Received: 30 April 2021; Accepted: 10 August 2021



(WAP) have shown the largest average atmospheric warming in the Southern Hemisphere, accompanied by significant warming of surface and deeper waters, salinity changes and rapid glaciers retreat (Henley *et al.*, 2019). Moreover, marine ecosystems will be affected depending on the duration and intensity of the heat wave (Frölicher and Laufkötter, 2018) and the capacity of marine biota to adapt their physiology to the exposure to altered conditions (Doney *et al.*, 2012).

In the marine realm, a number of environmental factors determine the productivity of phytoplankton assemblages, many of which are affected by global climate change, ozone depletion and pollution (Winder and Sommer, 2012; Häder and Gao, 2015; Häder *et al.*, 2015). Temperature is a crucial factor, which controls growth and productivity through phytoplankton photosynthesis (Thyssen *et al.*, 2011; Raymont, 2014; Hernando *et al.*, 2015). In this manner, productivity shows a positive correlation with the increase in temperature up to species optimum (Huertas *et al.*, 2011). Changes in species composition are already being observed in response to altered temperature patterns (Larsson *et al.*, 2015; Hernando *et al.*, 2015, 2020; Antoni *et al.*, 2020), considering that the ability to adapt to such changes are species-specific in a phytoplankton assemblage (Gao *et al.*, 2012). Consequently, the pronounced variability in ocean-atmosphere exchanges of heat along WAP has a strong impact on primary production, community composition and ecosystem functioning (Ducklow *et al.*, 2013; Meredith *et al.*, 2017; Hernando *et al.*, 2015).

In freshwater systems, high inputs of nutrients usually favor phytoplankton growth and photosynthesis (Falkowski *et al.*, 1998; Behrenfeld *et al.*, 2006), which promoted an increase in the frequency of cyanobacterial blooms in the last decades. Additionally, warmer temperatures and eutrophication are often proposed as principal factors favoring these events (Chorus and Bartram, 1999; Paerl and Huisman, 2008). These blooms are further associated with a decrease in phytoplankton diversity (Oliver and Gand, 2000). Other environmental factors promoting the predominance of cyanobacteria are irradiance conditions (with optimum values between spring and autumn), the ability to fix atmospheric nitrogen, high pH (6.5 to 8.5), low rate of filtration by zooplankton and the ability to form gas vesicles (Martin, 2000; Paerl and Huisman, 2008).

Cyanobacteria are an excellent source of peptides, FAs, amino acids, vitamins, minerals and pigments (Mimouni *et al.*, 2012). On the other hand, diatoms are the most abundant phytoplankton species and major primary producers in the oceans and are recognized as the most ecologically successful microalgae (Obata *et al.*, 2013). Diatom's membranes are usually enriched with medium-chain FAs as well as very long chain polyunsaturated fatty acids (PUFA) (Zulu *et al.*, 2018). These unsaturated FAs (UFAs) play vital roles in membrane physiology. In addition, the ratio of UFAs to saturated FAs (SFA) determines membrane fluidity, which promotes several cellular activities (Altabe *et al.*, 2013). The optimal membrane fluidity to sustain physiological homeostasis is achieved by upward or downward regulation of UFAs synthesis ('homeoviscous adaptation', Hazel, 1995). Temperature stress may induce changes in the FAs of cell membranes to avoid damage and be protected against the effects of increased

temperature (de la Rosa *et al.*, 2020). This adaptation allows cyanobacteria and microalgae such as diatoms to survive in extreme conditions and involves remodeling membrane lipids by modifying FAs chain length and unsaturation to sustain the desired level of fluidity in cell membranes (Sinensky, 1974).

Changes in PUFA in response to environmental stressors are highly relevant for the ecological role of cyanobacteria and microalgae as food sources for grazers and can therefore have serious implications for the flow of energy and the overall functioning of the ecosystem (Twining *et al.*, 2021).

The study of the changes induced by elevated seawater temperature in FAs and lipid metabolism enzymes of diatoms can provide insight into the impacts of marine heat waves on these organisms and on the potential repercussions on the entire marine food webs and ecosystems (Galloway and Winder, 2015). Changes in FAs metabolism and alterations in the expression of key genes (An *et al.*, 2013; Shimojima *et al.*, 2009) could be a response to environmental stresses such as heat waves.

Overall, organisms' responses to heat waves may involve reversible adjustment of their physiology by changing the content of some biomolecules or with activation of antioxidant systems, among other processes (Hernando *et al.*, 2015; Hernando *et al.*, 2018; Hernando *et al.*, 2020; de la Rosa *et al.*, 2020). Because phytoplankton growth and composition are highly dependent on environmental conditions (Galloway and Winder, 2015; Antoni *et al.*, 2020), there is an urgent need to understand how microalgae will be affected at the biochemical level in our warming world. Also, algae inhabiting regions at different latitudes are adapted to different ranges of temperatures.

The aim of this review is to compare the microalgae and cyanobacteria response from cold vs. temperate and tropical waters, to heat waves in relation to climate change and summarize the mechanisms of action leading to the observed FAs changes.

Importance of Fatty Acids and Their Role in Ecosystems

There are two major groups of FAs: SFA and UFAs, which differentiate from each other by the existence of double bonds in UFAs. The position of the double bonds is determined by the desaturase enzyme activity, which performs its activity in different positions accordingly to the type of enzyme present in different organisms (Brett and Müller-Navarra, 1997). SFA are mainly considered a source of energy and lipid storage. UFAs can be identified by the number of double bonds: Monounsaturated FAs (MUFAs) have one double bond and can be synthesized *de novo* almost by all organisms (Arts *et al.*, 2001). PUFA are FAs of 18 carbons or more in length with two or more double bonds. They can be classified into n-6 (or ω 6) and n-3 (or ω 3) major groups, depending on the position of the first double bond proximate to the methyl end of the FAs.

FAs synthesis in algae occurs through an aerobic pathway (Monroig *et al.*, 2013), it takes place in the chloroplast and the endoplasmic reticulum (ER). The glucose produced by photosynthesis is converted by glycolysis to pyruvate, which is the molecular basis for all metabolism processes. The desaturation and elongation process takes place in the ER. It uses 18:1 ω 9 to form linoleic

acid (LA; 18:2 ω 6) by Δ 12 desaturase and further into either the ω 6 pathway by Δ 6 desaturase or into the ω 3 pathway by Δ 15 desaturase to form alpha-linolenic acid (ALA; 18:3 ω 3). Different organisms utilize a diverse amount of desaturases to come to the different PUFA as shown in Fig. 1.

In consumers, PUFA play an essential role in brain development (Liu *et al.*, 2015) and many physiological functions such as down regulating inflammation and cellular signaling (Stillwell and Wassall, 2003). Essential FAs are some PUFA that play major important functions in physiological and biochemical processes. These must be acquired through dietary input, considering that the majority of the animals cannot synthesize them *de novo* (Kattner and Hagen, 2009) due to the lack of the desaturase enzyme (Sargent *et al.*, 1993).

Phytoplankton accounts for nearly 50% of net primary production on Earth, and is the source of many biomolecules (Field *et al.*, 1998) such as FAs (Guschina and Harwood, 2009) which are crucial components of the biosphere (Beardall *et al.*, 2009). PUFA play an important role in the food web because they affect key physiological processes and are precursors to many hormones (Jónasdóttir *et al.*, 2009; de Troch *et al.*, 2012). Thus, PUFA are critical regulators of the survival, reproduction, and

population growth in invertebrates and fish (Copeman *et al.*, 2002; Von Elert, 2004).

Long-chain ω 3 FAs, particularly eicosapentaenoic acid (EPA, 20:5 ω 3), docosapentaenoic acid (DPA, 22:5 ω 3), and docosahexaenoic acid (DHA, 22:6 ω 3) have been specially studied due to their important physiological functions and because the human body is unable to synthesize these compounds *de novo* (Drouin *et al.*, 2019). The brain has a high content of DHA, representing 50% of the total brain lipid and 10–15% of all of the FAs (Diau *et al.*, 2005). EPA was also found to be present at low concentrations in the brain (Chen *et al.*, 2009) probably because it rapidly converts to DHA (Kaur *et al.*, 2010). EPA in the brain provides important anti-inflammatory and immunity functions, but also plays an important role in neurotransmission and synaptic plasticity (Tassoni *et al.*, 2008; Uauy *et al.*, 2000).

In general terms, lipids produced by phytoplankton provide energy and essential nutrients for consumers in aquatic (Guschina and Harwood, 2009; Parrish, 2013) and terrestrial ecosystems (Gladyshev *et al.*, 2009; Hixson *et al.*, 2015). The herbivorous zooplankton and the fish that prey on them cannot synthesize all the FAs or cannot synthesize them in amounts required for optimal physiological performance (Brett *et al.*, 2006; Arts *et al.*, 2009). Hence, consumers acquire the required essential FAs from dietary lipids supplied by phytoplankton FAs, for example, LA, ALA, EPA, arachidonic acid (ARA, 20:4 ω 6) and DHA. In addition to their nutritive role, lipids form the backbone of the lipid bilayer in cell membranes (Arts *et al.*, 2009; Parrish, 2013). Although microalgae are the primary producers of EPA and DHA in marine food webs, humans use marine fish oil as the principal source of these PUFA (Sinclair, 2000). Moreover, the marine fish industry is increasingly declining due to overfishing and environmental pollution, while the production costs of DHA or EPA from cultured algae are potentially equal to the cost of producing EPA from fish oil (Milledge, 2011) and lower than the costs of PUFA from shellfish flesh and by-products (Al Khawli *et al.*, 2019; Dvoretzky *et al.*, 2021). Different algal taxa exhibit different FAs profiles (Harwood, 1998; Napolitano, 1999) as shown in Table 1 and Fig. 2. Jónasdóttir (2019) revealed a phyla-specific, and a highly species-specific PUFA production of marine phytoplankton, showing that the highest proportion of PUFA is found in Chlorophyta and Cryptophyta, and the lowest in Ochrophyta, Cyanobacteria and diatoms (Maltse and Maltseva, 2021). Diatoms are an important nutrition source in the marine environment, since many species produce large quantities of EPA (Arao *et al.*, 1987; Dunstan *et al.*, 1994). Both EPA and DHA are particularly found in diatoms, dinoflagellates and prymnesiophytes (Tonon *et al.*, 2002; Mansour *et al.*, 2005), they are produced from their precursor, ALA which is considered an essential FAs (Cook and McMaster, 2004).

Some ω 3 are mainly synthesized by producers in aquatic environments (Hixson *et al.*, 2015; Twining *et al.*, 2015). These PUFA are progressively consumed and selectively retained by other aquatic organisms higher up in the food chain (Kainz *et al.*, 2004; Hixson *et al.*, 2015), and transferred to terrestrial ecosystems via consumption (Gladyshev *et al.*, 2009, 2013). Thus, PUFA composition of




FIGURE 1. Simplified desaturation pathway of the FAs synthesis in phytoplankton cells. Des: Desaturase, Elo: elongase. Modified from Jónasdóttir *et al.* (2009) and Vaezi *et al.* (2013). The arrows indicate potential changes in the ω 6 and ω 3 FAs production path as a function of temperature. Red and black arrows indicate high or low temperature conditions, respectively.

TABLE 1
FAs composition in different edible marine and freshwater sources in cold and temperate environments

| Region | Phytoplankton group/Species | Exposure conditions (Time/Stressor) | SFA | MUFA | PUFA | UFA/SFA | ω 3 content at the end of experiments | Reference |
|-----------|--|---|------------------|---|---|----------------|--|-------------------------|
| Temperate | <i>Microcystis aeruginosa</i> (A) | 7 days/T+ | 16:0; 18:0 | cis-9-18:1 | GLA; ALA | N/ch | ↗ | de la Rosa et al., 2020 |
| Antartic | <i>Chlamydomonas</i> sp (B) | 56 hs/T (-20, -10, 0, 5°C) | 16:0; 18:0 | 16:1 | 18:3; 20:5; 20:3 | N/ch | ↖ | An et al., 2013 |
| Antartic | Diatoms, Prasinophytes, Cryptophytes, Prymnesiophytes and others (C) | 6 days/S-T+; S0T+; S-T0 | 16:0; 14:0; 18:0 | 16:1 ω 7 | 16:3 ω 3; SDA; EPA; DHA | ↗ (S-T+; S0T+) | ↖ (S-T+; S0T+; S-T0) | Hernando et al., 2018 |
| Temperate | Diatoms and Green algae | T+ | N/A | N/A | ALA; EPA; DHA; LA; ARA | N/A | ↘ | Arts et al., 2015 |
| Temperate | <i>Thalassiosira pseudonana</i> (D) | 500 generations/T+ (16, 31°C) | 16:0; 18:0 | 16:1 Δ 3; 16:1 Δ 7 | EPA; DHA; 16:3 ω 3; 16:4 ω 3; ALA | N/A | N/A | O'Donnell et al., 2019 |
| Temperate | <i>Thalassiosira weissflogii</i> (E) | 7 days/S0T+; S-T+; S+T+; S0T-; S+T-; S-T-; S0 | 23:0; 21:0; 20:0 | 16:1; 17:1 | 20:3 ω 3; ARA; EPA; DHA | ↘ (S0T+; S+T+) | ↗ (S0T+; S+T+) | Gonçalves et al., 2017 |
| Temperate | <i>Scenedesmus obliquus</i> (F) | 2 weeks/T+ | 16:0; 18:0 | 18:1 ω 9 | LA; ALA; SDA | ↘ | ↗ | Fuschino et al., 2011 |
| Antartic | Coastal phytoplankton assemblage (C) | 7 days/S0T0; S-T0; S0T+; S-T+ | 14:0; 16:0 | 16:1 ω 7 | SDA; EPA | ↖ (S0T+; S-T+) | ↗ (S0T+; S-T+) | Antacil et al., 2021 |
| Temperate | <i>Nannochloropsis</i> sp. (G) | 7 days/L+S+, L+S0, L+S-, L-S+, L-S0, L-S- | 16:0; 14:0 | 16:1; 18:1 ω 9 | LA; ARA; EPA | N/A | ↖ low salinity, independent of light intensity | Pal et al., 2011 |
| Polar | <i>Chaetoceros brevis</i> , <i>Pyramimonas</i> sp. (H) | L-T+; L+T+; L-T-; L+T- | 16:0 | 16:1 ω 7 (except <i>Pyramimonas</i> sp.) | SDA; 16:4 (EPA only in <i>C. brevis</i> and DHA only in <i>Pyramimonas</i> sp.) | N/A | ↖ | Boelen et al., 2013 |
| Temperate | <i>Thalassiosira weissflogii</i> , <i>Emiliania hux</i> and <i>Fibrocapsa japonica</i> | L-T+; L+T+; L-T0; L+T0 | 16:0; 14:0 | 16:1 ω 7 (except <i>E. hux</i>) | SDA (EPA except <i>E. hux</i> and DHA except <i>F. japonica</i>) | N/A | ↘ | |

(Continued)

Table 1 (continued).

| Region | Phytoplankton group/Species | Exposure conditions (Time/Stressor) | SFA | MUFA | PUFA | UFA/SFA | ω 3 content at the end of experiments | Reference |
|--------------|--|---|------------------------------------|---|--|---------|--|------------------------------------|
| Temperate | <i>Phaeodactylum tricornutum</i> (I), <i>Chaetoceros muelleri</i> (I) | 2 h/T+ (30, 35, 40°C) and 24 h/T+ (30, 35°C) | 16:0; 14:0 | 16:1 | EPA | ➔ | ➔ | Rousch <i>et al.</i> , 2003 |
| Temperate | <i>Phaeodactylum tricornutum</i> (I) | 5 days/Heat waves (18, 26°C) | 16:0; 14:0 | 16:1 | 18:2 ω 6; 18:3 ω 6; 20:3 ω 6; EPA; 18:4 ω 3; 16:3 ω 4 | ➔ | ➔ | Feijão <i>et al.</i> , 2020 |
| Antarctic | Sea-ice diatom <i>Nitzschia lecointei</i> (K) | 7 days/S-T+ (-1.8; 3°C) | 16:0; 14:0; 18:0 | 16:1 ω 7; 18:1 ω 9 | 18:3 ω 3; 18:4 ω 3; EPA; 22:6 ω 3 | ↔ | ↔ | Torstensson <i>et al.</i> , 2019 |
| Temperate | <i>Odontella aurita</i> (L) | 10 days/8, 16, 24°C | 14:0; 16:0; 18:0 | 16:1 ω 7; 18:1 ω 9 | 18:4 ω 3; 20:4 ω 6; EPA; 22:6 ω 3 | ➔ | ➔ | Pasquet <i>et al.</i> , 2014 |
| Tropical | <i>Chaetoceros sp.</i> ; <i>Rhodomonas sp.</i> ; <i>unidentified Prymnesiophyte</i> (M) | Exposed to 25, 27, 30 and 33°C. Harvested in late logarithmic grown phase | 12:0; 14:0; 16:0; 17:0; 18:0 | 16:1 ω 7; 18:1 ω 9 18:1 ω 7 | 16:2 ω 7; 18:2 ω 6; 18:3 ω 6; EPA; 18:4 ω 3; 22:6 ω 3 | ➔ | ➔ | Renaud <i>et al.</i> , 2002 |
| Sub-tropical | <i>Nannochloropsis salina</i> (N) | 14 days/17, 21 and 26°C | 14:0; 16:0 | 16:1; 18:1 ω 9 | 20:4 ω 6; 18:2 ω 6; 18:3 ω 6; EPA | ➔ | ➔ | Hoffmann <i>et al.</i> , 2010 |
| Polar | <i>Leptolyngbya sp.</i> (O) | Ambient conditions | 18:0; 12:0; 17:0 | 16:1 ω 9; 18:1 ω 7 | 18:2 ω 6; 18:3 ω 6; 18:3 ω 3; EPA | N/A | N/A | Zainal Abidin <i>et al.</i> , 2020 |

Note: The effect of different treatments (increased temperature (T+), decreased salinity (S-), light conditions) are shown. N/A: not available; N/Ch: not significant changes; T: temperature; S: salinity; L: light intensity; 0: control; WT: wild type; ARA: arachidonic acid, 20:4 ω 6; GLA: gamma linolenic acid, 18:3 ω 6; ALA: alpha-linolenic acid, 18:3 ω 3; SDA: stearidonic acid, 18:4 ω 3; EPA: eicosapentaenoic acid, 20:5 ω 3; DHA: docosahexaenoic acid, 22:6 ω 3; LA: linoleic acid, 18:2 ω 6; POA: palmitoleic acid, 16:1 Δ 9.




FIGURE 2. Distribution of studies on the effects of temperature increase on microalgae and cyanobacteria FAs composition. Red dots denote positioning and letters from the Roman alphabet correspond to the phytoplankton species or groups indicated in Table 1.

phytoplankton is an important determinant of food quality and, consequently, the health and optimal functioning of both aquatic and terrestrial consumers (Budge *et al.*, 2002; Dalsgaard *et al.*, 2003).

Changes in Membrane Fatty Acids

Temperature has strong directional effects on both the quantity and quality of FAs in phytoplankton. Phytoplankton and other organisms adapt to changing temperatures by modifying the structure of their membranes (Winter and Dzwolak, 2005), a process known as homeoviscous adaptation (see above). The double bonds in PUFA increase flexibility, leading to increased membrane fluidity. Under a sudden change in environmental conditions, *de novo* synthesis of UFAs cannot occur rapidly, but the desaturation of FAs may be adjusted by transferring specific acyl groups to other polar lipids and allowing rapid adaptive membrane reorganization (Khozin-Goldberg and Cohen, 2006). Temperature changes may then induce the replacement of an enzyme by an isoenzyme with better heat- or cold-tolerance (Steele and Fry, 2000). Several authors have shown an inverse relationship between temperature and FAs unsaturation in microalgae (Wada *et al.*, 1990; Zhu *et al.*, 1997).

PUFA represent between 30 and 60% of total FAs content in natural polar ocean assemblages and in individual phytoplankton species (Teoh *et al.*, 2004). Hernando *et al.* (2018) described the predominance of PUFA (44–64% in average) over SFA for Antarctic coastal phytoplankton at the beginning of microcosm experiments to evaluate the effects of increased temperature and decreased salinity on physiology (Table 1, Fig. 2).

In polar region, cyanobacteria mats from streams and lakes are dominated by Nostoc or oscillatoriaceae (Oscillatoriaceae) (Zainal Abidin *et al.*, 2020) with extreme accumulations up to 90 cm thick and $> 40 \mu\text{g Chla cm}^{-2}$ at some sites. Picocyanobacteria often dominate the phytoplankton in polar lakes achieving some of the highest natural concentrations on record, up to 8×10^6 cells mL^{-1} in some Antarctic lakes

(Vincent, 2000). In marine waters in the Southern Ocean, picocyanobacteria were found to decrease during spring (Massana *et al.*, 1998) and summer (Murray *et al.*, 1998), and it has been speculated that the significant decline of archaea during spring/summer is a result of the competition with non-archaeal microorganisms during phytoplankton blooms (Massana *et al.*, 1998). Regarding FA studies in these organisms, Zainal Abidin *et al.* (2020) demonstrated a high content of UFAs in four of the five isolated species of the filamentous cyanobacterium *Leptolyngbya* sp. from fine Antarctic mats. The $\omega 6$ linoleic acid reached an average percentage of 60% followed by palmitoleic acid (10%) and oleic acid (10%) as shown in Table 1 and Fig. 2.

Contrarily to observations in Antarctic waters, Ohse *et al.* (2015) found in a FAs study with temperate marine species (*Nannochloropsis oculata*, *Thalassiosira pseudonana*, *Phaeodactylum tricorutum*, *Isochrysis galbana*, *Tetraselmis suecica*, *Tetraselmis chunii*, *Chaetoceros muelleri*, *Thalassiosira fluviatilis* and *Isochrysis* sp.) and the freshwater *Chlorella* sp, that average PUFA in these species was around 20%. Also, the cyanobacteria *Microcystis aeruginosa*, showed a PUFA relative abundance around 40% of the total FAs (de la Rosa *et al.*, 2020).

Water temperature is one of the most important factors in the regulation of PUFA synthesis by microalgae or cyanobacteria and their accumulation by zooplankton. Thus, global warming may lead to a decrease of PUFA in aquatic ecosystem (Thompson *et al.*, 1992; Renaud *et al.*, 2002; Maazouzi *et al.*, 2008) in order to maintain fluidity (see above). Conversely, acclimation to increasing temperature involves decreasing PUFA membrane content, while, simultaneously, increasing SFA (Rousch *et al.*, 2003; Fuschino *et al.*, 2011) as shown in Table 1.

Consequences of the Increase in Temperature in the Composition of FAS in Temperate and Tropical vs. Cold Environments

An important consequence of climate change is that increasing water temperatures are expected to reduce the

global production of PUFA in phytoplankton. The environment temperature and salinity affect the FA profile of microalgae, and it was found that DHA content decreases significantly with an increase in water temperature and salinity (Adarme-Vega *et al.*, 2012). Similarly, the EPA content was also found to decrease considerably with an increase in water temperature (Sang *et al.*, 2012). Changes in the biochemical composition of phytoplankton cell membranes may lead to cascading effects throughout the world's aquatic ecosystems. Further, such modified biochemical composition in aquatic ecosystems, as mentioned above, is also anticipated to propagate to terrestrial animals because of the flux of aquatic biomass, containing ω 3 PUFA (Gladyshev *et al.*, 2009; Gladyshev *et al.*, 2013). This is critical because PUFA not only enhance the growth rates and reproductive capacities of aquatic animals (Ballantyne *et al.*, 2003), but are also vitally important to the neural/cognitive, cardiovascular, and visual health of terrestrial vertebrates (Calder, 2015).

Antarctic Waters

Rapid changes in temperature and salinity may have great effects on the physiology and biochemical composition of Antarctic diatoms. Hernando *et al.* (2018) and Antacli *et al.* (2021) demonstrated that EPA increased in Antarctic coastal phytoplankton assemblage in response to stress (increased temperature combined with freshening), in concordance with results from Teoh *et al.* (2013) as shown in Table 1 and Fig. 2. An *et al.* (2013) working with the Antarctic ice algae *Chlamydomonas* sp. and considering the FA desaturase (FADs) expression, demonstrated an increased amount of EPA in cells exposed to 0 and 5°C in comparison to controls at -10°C. This is related to the strategies adopted for survival at low temperatures, which include an increasing degree of FAs unsaturation or branching and a decrease in the FAs chain length (Penfield, 2008). An *et al.* (2013) results indicated that the mRNAs of FADs in *Chlamydomonas* sp. were regulated differentially in response to temperature stress. Moreover, those desaturases involved in ω 3 production as shown in Fig. 1 were found to be sensitive to low temperature, indicating that they might play primary roles in protecting algal cells from potential damage at low temperature. Furthermore, when cells of the Antarctic sea-ice diatom *Nitzschia lecointei* were grown at 3°C, they had higher relative PUFA content (mainly hexadecatetraenoic [16:4 ω 4] and EPA) and lower relative MUFAs content than cells grown at -1.8°C (Torstensson *et al.*, 2019). Diatoms isolated from sea ice show a FAs composition characterized by high levels of EPA and C16 PUFA (Henderson *et al.*, 1998). Analysis of the FAs of a natural Arctic phytoplankton assemblage showed that, when dominated by diatoms of the genera *Thalassiosira* and *Chaetoceros*, the proportions of C16:1(ω 7) and EPA were correspondingly high (Reuss and Poulsen, 2002).

Most algae produce EPA via the ω 3 pathway as shown in Fig. 1. The biosynthesis of EPA occurs through a series of reactions that can be divided into two distinct steps. First is the *de novo* synthesis of oleic acid (OA, 18:1 ω 9) from acetate. OA is further desaturated by a Δ 12 desaturase to

form LA and a Δ 15 desaturase to form ALA. Then, ω 6 and ω 3 FAs families are formed from their precursors by a series of desaturation and elongation reactions, including EPA as shown in Fig. 1. The three FAs, OA, LA and ALA, compete with each other for the Δ 6 desaturase. The affinity of the enzyme to the substrate and the amount of substrate available determine which metabolic pathway is predominant (Gurr, 1985). Generally, the first Δ 6 desaturation is the limiting step and ALA has the highest affinity for Δ 6 desaturase followed by LA and oleic acid (Jónasdóttir, 2019). In order to enhance EPA production in *Dunaliella salina* (a green alga), FAs desaturases (Δ 6) FADS6 from the diatom *Thalassiosira pseudonana*, Shi *et al.* (2018) constructed and overexpressed in *D. salina* with a plasmid (Ds-TpFADS6). They demonstrated that EPA production increased significantly in *D. salina* transformants. However, after Ds-TpFADS6 transformants were maintained at low temperature (4°C) for 5 days, their ALA levels increased but LA levels decreased. From this, they concluded that the activity of other desaturase enzymes overexpressed in *D. salina*, Ds-FADS15 (Δ 15), was enhanced at low temperature, promoting the conversion of LA to ALA and increasing the ALA level as show in Fig. 1. These findings are similar to those of Okuda *et al.* (2015), who observed an accumulation of EPA in *Mortierella alpina* at low temperature (below 15°C). Ds-FADS15 and Ds-FADS6 in *M. alpina* transformants had high activity and exhibited greater EPA accumulation under varying temperatures than under constant temperatures, suggesting a more active LA-ALA-EPA pathway under these conditions. There was also less LA flux through the ω 6 pathway as shown in Fig. 1.

EPA or other long-chain PUFA (LC-PUFA) may play a role in the functioning of the thylakoid membrane and are therefore essential for photosynthesis (Cohen *et al.*, 1988). At higher irradiance, photosynthesis is less efficient and thus fewer thylakoid membranes are required. As a result, LC-PUFA content could be lower in high light-acclimated algae (Harwood and Jones, 1989). Hence, cold-adapted polar microalgae could have high EPA and DHA production, since they can exhibit high growth rates under low temperature and irradiance conditions. Furthermore, polar species are expected to sustain a higher EPA and DHA content, since their habitat is characterized by lower average irradiance and temperature levels (Schloss *et al.*, 2002).

Changes in salinity may also modify the composition of FA and influence the physiological properties of microalgae (Hernando *et al.*, 2018). Due to the increase in the occurrence of heat waves in the WAP, melt water and seawater freshening increased. Xu and Beardall (1997) demonstrated that the content of ω 3 in *Dunaliella* sp. decreased as the average salinity increased. In a culture of the marine diatom *Nitzschia laevis*, EPA yield was the highest at half the salinity of the artificial seawater (Wen and Chen, 2001). Similarly, both EPA and 18:4 ω 3 were significantly higher in natural Antarctic phytoplankton assemblages after 48 h exposure at salinities 4 PSU lower than environmental values (Hernando *et al.*, 2018).

High level of intracellular molecular oxygen may be another reason for the increased PUFA at low temperature, because the enzymes responsible for desaturation and elongation of PUFA depend on the availability of molecular

oxygen (Chen and Johns, 1991; Singh and Ward, 1997). A temperature shift strategy has been employed to enhance the overall ω 3 PUFA production (including EPA) because the optimal temperature for microalgal growth is often higher than that for ω 3 PUFA formation (Jiang and Chen, 2000). Such a phenomenon has been observed in many different algal species including *Porphyridium cruentum* (Springer et al., 1994), *Nannochloropsis* sp. (Sukenik, 1991) and *P. irregulare* (Stinson et al., 1991).

To the best of our knowledge, there are no studies testing the effects of temperature increase on FA of Antarctic cyanobacteria. In a recent latitudinal study, Breton et al. (2020) demonstrated that the distribution of marine *Synechococcus* cyanobacteria depends on the differentiation of lineages adapted to distinct thermal environments. Over the temperature gradient they sampled, all strains maintained efficient photosynthetic capacities. Subpolar and cold temperature strains actually showed enhanced capacities for lipid monodesaturation thanks to an additional Δ 9-desaturase. By contrast, tropical and warm temperature strains displayed moderate monodesaturation capacities but high proportions of double unsaturations in response to cold temperatures, thanks to Δ 12-desaturases. The desaturase genes displayed specific distributions directly related to latitudinal variations in ocean surface temperature. Thus, this study highlights the critical importance of membrane fluidity modulation by desaturases in the adaptive strategies.

Temperate and Tropical Waters

Many studies have shown changes in the FAs composition of microalgae and cyanobacteria from temperate waters when exposed to changes in temperature. *Phaeodactylum tricornerutum* has been used as a model organism for diatom physiology studies, being the first pennate diatom to have its genome completely sequenced (Bowler et al., 2008). It has been shown that this species has the ability to produce up to 35% of EPA of total FA (Hamilton et al., 2014). The thylakoid membrane of *P. tricornerutum* is composed of uncharged glycolipids such as monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG) (Yang et al., 2017), with EPA being the most abundant component of the glycolipids (Mühlroth et al., 2013). Jiang and Gao (2004) explained that elevated levels of UFAs in the diatom *P. tricornerutum* after 12 h of exposure to low temperatures were due to increased activity of desaturase. Other studies have shown similar changes in the FA composition in diatoms within a similar period of time (12–48 h) (Tonon et al., 2002; Siron et al., 1989; Liang et al., 2006). Thompson et al. (1992) who studied the effect of temperature over the range from 10 to 25°C on the FA composition of eight species of marine phytoplankton, including *Thalassiosira pseudonana* and three species within the genus *Chaetoceros* also showed a trend towards elevated PUFA at a lower temperature. However, only for *T. pseudonana* the percentage of DHA decreased linearly with increasing temperature. Blanchemain and Grizeau (1999) noted that the diatom *Skeletonema costatum* increased its EPA levels relative to dry weight at only the lowest (15°C) of three treatment temperatures and that

there was a positive correlation between treatment duration (4 or 15 h) and EPA production at a particular temperature.

Species of the genus *Nannochloropsis* (Eustigmatophyceae, Monodopsidaceae) from sub-tropical sea waters are an important source of the essential EPA and are widely used in marine aquaculture nutrition (Renaud and Parry, 1994). The quality of *Nannochloropsis* biomass for aquaculture (high EPA percentage of total FAs) has been shown to deteriorate with culture age and with increasing light intensity, salinity level, and temperature. In addition, a concomitant increase in the proportions of saturated and monounsaturated fatty acids 14:0, 16:0, and 16:1 was found (Sukenik et al., 1993; Renaud and Parry, 1994; Roncarati et al., 2004). Hoffmann et al. (2010) also described this trend for *Nannochloropsis salina* with increased EPA contents under a low culture temperature (17°C) as compared to moderate temperatures of 21°C and 26°C (Table 1, Fig. 2).

Renaud et al. (2002) also demonstrated a decrease in the total production of the PUFA, including significant decreases in the percentages of the 20:5 ω 3 and 22:6 ω 3 in three tropical Australian species of *Rhodomonas* sp., an unidentified prymnesiophyte and a *Chaetoceros* sp. at higher growth temperatures between 33°C and 35°C when harvested during in late logarithmic growth phase. These results concurs with the findings of previous studies (Mortensen et al., 1988; Renaud et al., 1995) as shown in Table 1 and Fig. 2.

For the temperate *Chlamydomonas*, temperature stress (variation from optimum) caused an increase in SFA (Teoh et al., 2013). The increase in SFA could be an important energy source for stress adaptation in these microalgae. The role of UFAs in membrane fluidity has been demonstrated in other algae such as *Spirulina platensis* (Sushchik et al., 2003), *Chlamydomonas* sp. (Poerschmann et al., 2004), and *C. vulgaris* strain BI (Morgan-Kiss et al., 2008) when subjected to temperature stress. Pal et al. (2011) demonstrated that the eustigmatophyte *Nannochloropsis* sp. accumulate energy-rich storage products when exposed to unfavorable conditions that limit growth as long as the carbon supply exceeds the photosynthetic capacity of the cells. Also, low temperatures resulted in higher lipid production in *Nitzschia palea* (Renaud et al., 1995), *Chaetoceros* sp. (Clone CS256) (Renaud et al., 2002) and *Chaetoceros* cf. *wighamii* (de Castro Araujo and Garcia, 2005).

Qiao et al. (2016) found a higher DHA/EPA ratio in *P. tricornerutum* grown under altered environmental conditions. DHA percentage increased with increasing light intensity, but decreased with increasing temperature. Furthermore, Feijão et al. (2017) also working with *P. tricornerutum* from temperate ocean waters, demonstrated that cells exposed to heat wave conditions showed lower relative EPA contents and double bond indexes, whereas the ω 6/ ω 3 ratio increased like in the cyanobacteria *Microcystis aeruginosa* (de la Rosa et al., 2020). Moreover, the analysis of the FAs profiles in *P. tricornerutum* also suggested that heat exposure negatively impacted thylakoid lipids, in agreement with the decrease observed in photosynthesis. Oxygen-evolving complexes are essential for the water splitting reaction occurring at the donor side of the photosystem II (PSII) and essential not only for oxygen production but also for fueling the quinone pool with the necessary energy for electron transport from

PSII to PSI (Strasser *et al.*, 2000). These complexes are usually heat-sensitive (Duarte *et al.*, 2016) and therefore heat waves may have a negative impact on photosynthesis and cells' nutritional value, as well as on their capacity to oxygenate ocean water.

In order to get further insights on the influence of gene expression in the changes of lipid classes and FAs composition on *P. tricornutum* as a consequence of heat waves, Feijão *et al.* (2020) analysed key genes responsible for the biosynthesis of some lipids, FAs desaturation of plastidial $\Delta 6$ desaturase and extraplastidial $\Delta 5$ desaturase lipids, as well as phospholipases. The results obtained indicate that storage lipids decrease in response to a heat wave. Besides changing the proportion of membrane/storage lipids, the heat wave also affected the FAs composition of all lipid classes. As a general trend, there is a decrease in PUFA and an increase in saturated or, in some cases, monounsaturated FAs. This could be a mechanism to stabilize the membrane (Feijão *et al.*, 2020). Moreover, Feijão showed that after exposure to heat, the gene encoding the plastidial $\Delta 6$ FAs desaturase showed a significant increase. Thus, although no significant change occurred in the $\Delta 5$ desaturase transcripts amounts, lower amounts of EPA were detected. Under extreme temperature stress (up to 30°C) *P. tricornutum* hardly shows any growth (Jiang and Gao, 2004) and effects on lipid metabolism are drastic (e.g. no EPA content) (Dodson *et al.*, 2014). This mechanism could involve gene expression regulation of genes related to membrane lipids synthesis e.g. MGDG (with high EPA content). Under non-lethal growth temperatures, *P. tricornutum* could prioritize thylakoid membrane stability and therefore photosynthesis over membrane quality (Feijão *et al.*, 2020).

The marine picocyanobacterium *Synechococcus* is one of the most conspicuous and widely distributed species in the world, with high importance for net primary productivity (Flombaum *et al.*, 2013). Studies of the FAs desaturase-related genes in *Synechococcus* have determined the presence of four main distinct genes, encoding two $\Delta 9$ - (desC3 and desC4) and two $\Delta 12$ -desaturases (desA2 and desA3; Varkey *et al.*, 2016). *Synechococcus* has a different physiological strategies based on the presence of specific lipid desaturase gene sets, allowing them to cope with changes in the fluidity of the photosynthetic membrane in different thermal conditions. In a recent latitudinal study, Breton *et al.* (2020) have shown the distribution of such lipid desaturase genes at 33 widespread stations along the Tara Oceans transect. The desaturase gene desC4, which is specific for cold-adapted strains, was found exclusively in waters colder than 20°C, consistent with its hypothesized role of a monodesaturation enhancer enabling growth in cold thermal environments. In warm environments, there is little need to have FAs variations in membrane, and marine *Synechococcus* have adopted a different strategy. They constitutively contain membrane $\Delta 12$ -lipids with more double unsaturation thanks to DesA2, a desaturase specific to warm environments (30–32°C). $\Delta 9$ -desaturases were selected in colder environments during the evolution of marine *Synechococcus* as shown in Fig. 1. Thus, after shifting the culture temperature down from 38 to 22°C, Sakamoto *et al.* (1997) demonstrated for *Synechococcus* sp. PCC 7002 strain that the degree of unsaturation of C18 FAs

increased in cells grown at low temperature. The desaturation at the $\omega 3$ position of C18 increased gradually during a 12-h period in cells grown at 22°C. Various desaturases in cyanobacteria and higher plants are known to be affected by temperature at the level of gene transcription (Los *et al.*, 1997). Wada and Murata (1990) demonstrated that FAs desaturases enzymes play an important role in cold acclimation of cyanobacteria. The microalgal genome contains genes coding for the synthesis of molecules involved in survival mechanisms and FA synthesis (Li *et al.*, 2014). In these organisms, de la Rosa *et al.* (2020) demonstrated a differential sensitivity of $\omega 3$ and $\omega 6$ FA to high temperature. They observed decreased relative concentration of 18:4 $\omega 3$ and 18:3 $\omega 3$ at 29°C after 10-days exposure as compared to exposure to 26°C. Previous studies also have shown an effect of temperature on the FA composition, and specifically a decrease in $\omega 3$ PUFA (EPA and DHA) with increasing temperature. Thus, Sushchik *et al.* (2003) demonstrated a decrease of $\omega 3$ FA desaturase due to increased temperature in *Chlorella vulgaris* and *Botryococcus braunii* resulted in a decrease in the relative content of the more unsaturated intracellular FAs, especially the trienoic FAs like ALA. Also, for *Thalassiosira pseudonana*, the percentage of 22:6 $\omega 3$ decreased linearly with increasing temperature over the range from 10 to 25°C (Thompson *et al.*, 1992).

Hence, the adaptation of *M. aeruginosa* to temperature rise could be related to a decrease and an increase in the activity of $\omega 3$ and $\omega 6$ -desaturases, respectively. The high relative abundance of cis-18:1 $\omega 9$, observed in high temperature conditions from day 1 of incubation and until the end of the 10-days experiment (de la Rosa *et al.*, 2020), could explain the significant increase of the PUFA $\omega 6$, considering that this MUFAs cis-18:1 $\omega 9$ is a precursor of the $\omega 6$ FAs (Akoh and Min, 2008).

Conclusions

FAs in phytoplankton and cyanobacteria display structural changes and high biological specificity, responding to stress conditions. Their biosynthesis is inhibited or enhanced depending on the temperature the organisms are exposed to, along with alterations in the processes of FAs desaturation and elongations. Hence, FAs are useful bioindicators of the ecological and health status of aquatic ecosystems. They provide crucial information about the impact of global stressors on aquatic communities and thus on the food web, with severe repercussions to human beings among other terrestrial animals, and food quality. Moreover, lipids and proteins are involved in the most vital functions of aquatic organisms. Thus, the reduction of essential FAs content at the base of the food web may have serious implications for higher trophic levels.

As consequences of climate change effects on marine and freshwater ecosystems, intensive agriculture production with the usage of fertilizers and pesticides near coastal wetlands will have severe impacts on the aquatic communities and thus to whole the ecosystem. Therefore, it is of major importance and becomes a priority to determine and predict the effects of environmental and anthropogenic stressors on the aquatic systems in order to promote their health and preserve biodiversity and food quality.

Overall, heat wave-induced changes in the relative abundance of PUFA, and more importantly EPA, are alarming because of their importance in aquatic ecosystems. However, the impact of the exposure of aquatic communities to climate change variables is different between temperate/tropical and polar environments. In cold marine waters the activation of desaturase is promoted towards high-energy ω 3 FAs such as EPA. Conversely, in temperate and tropical environments, the decrease in EPA as a function of desaturases sensitivity protects the photosystems in conditions of high irradiance and heat waves.

Availability of Data and Materials: Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Author Contributions Statement: The authors confirm contribution to the paper as follows: MH and IS participated in the designing of the original idea and in the manuscript writing. FD and MD contributed to the bibliographic search and in the manuscript writing. All authors have approved the final version of the article.

Funding Statement: The research leading to these results further received funding grants ANPCYT PICT 2011-130 Raíces of the Agencia Nacional de Promociones Científicas of Argentina to IRS and MPH, and from the European Union's Horizon 2020 research and innovation program under grant agreement No. 730984, ASSEMBLEPLUS2019359 Project awarded to FD. It is further a contribution to CoastCarb (Funding ID 872609, H2020, MSCA-RISE-2019, Research and Innovation Staff Exchange).

Conflicts of Interest: The authors declare that they have no conflicts of interest to report regarding the present study.

References

- Abhilash C (2015). Carbon footprint—A review. *International Journal of Scientific Research Today* **2**: 116–126.
- Adarme-Vega TC, Lim DK, Timmins M, Vernen F, Li Y, Schenk PM (2012). Microalgal biofactories: A promising approach towards sustainable n-3 fatty acid production. *Microbial Cell Factories* **11**: 96–106.
- Akoh CC, Min DB (2008). Food lipids: chemistry, nutrition, and biotechnology. In: Akoh CC, Min DB (eds.), *Food Lipids: Chemistry, Nutrition, and Biotechnology*, Third edition, pp. 427–432. CRC Press, Taylor and Francis Group, New York.
- Al Khawli F, Pateiro M, Domínguez R, Lorenzo JM, Gullón P, Kousoulaki K, Ferrer E, Berrada H, Barba FJ (2019). Innovative green technologies of intensification for valorization of seafood and their by-products. *Marine Drugs* **17**: 689.
- Altabe SG, Mansilla MC, de Mendoza D (2013). Remodeling of membrane phospholipids by bacterial desaturases. In: Ntambi J (ed.), *Stearoyl-CoA Desaturase Genes in Lipid Metabolism*, pp. 209–231. New York: Springer.
- An M, Mou S, Zhang X, Ye N, Zheng Z, Cao S, Xu D, Fan X, Wang Y, Miao J (2013). Temperature regulates fatty acid desaturases at a transcriptional level and modulates the fatty acid profile in the Antarctic microalga *Chlamydomonas* sp. ICE-L. *Bioresource Technology* **134**: 151–157.
- Antacli JC, Hernando MP, de Troch M, Malanga G, Mendiolar M, Hernández DR, Varela DE, Antoni J, Sahade RJ, Schloss IR (2021). Ocean warming and freshening effects on lipid metabolism in coastal Antarctic phytoplankton assemblages dominated by sub-Antarctic species. *Science of the Total Environment* **790**: 147879.
- Antoni JS, Almandoz GO, Ferrario ME, Hernando MP, Varela DE, Rozema PD, Buma AGJ, Paparazzo FE, Schloss IR (2020). Response of a natural Antarctic phytoplankton assemblage to changes in temperature and salinity. *Journal of Experimental Marine Biology and Ecology* **532**: 151444.
- Arao T, Kawaguchi A, Yamada M (1987). Positional distribution of fatty acids in lipids of the marine diatom *Phaeodactylum tricornutum*. *Phytochemistry* **26**: 2573–2576.
- Arts MT, Ackman RG, Holub BJ (2001). Essential fatty acids in aquatic ecosystems: A crucial link between diet and human health and evolution. *Canadian Journal of Fisheries and Aquatic Sciences* **58**: 122–137.
- Arts MT, Brett MT, Kainz MJ (2009). *Lipids in Aquatic Ecosystems*. New York: Springer, 308.
- Arts MT, Hixson SM, Mercieca S (2015). Climate warming reduces essential fatty acid production in algae. *Journal of Environmental Indicators* **9**: 7–8.
- Ballantyne AP, Brett MT, Schindler DE (2003). The importance of dietary phosphorus and highly unsaturated FAs for sockeye (*Oncorhynchus nerka*) growth in Lake Washington—A bioenergetics approach. *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 12–22.
- Beardall J, Ihnken S, Quigg A (2009). Gross and net primary production: Closing the gap between concepts and measurements. *Aquatic Microbial Ecology* **56**: 113–122.
- Behrenfeld M, O'Malley R, Siegel D, McClain CR, Sarmiento JL, Feldman GC, Milligan AJ, Falkowski PG, Letelier RM, Boss ES (2006). Climate-driven trends in contemporary ocean productivity. *Nature* **444**: 752–755.
- Blanchemain A, Grizeau D (1999). Increased production of eicosapentaenoic acid by *Skeletonema costatum* cells after decantation at low temperature. *Biotechnology Techniques* **13**: 497–501.
- Boelen P, van Dijk R, Damsté JSS, Rijpstra WIC, Buma AG (2013). On the potential application of polar and temperate marine microalgae for EPA and DHA production. *AMB Express* **3**: 1–9.
- Bowler C, Allen A, Badger J, Grimwood J, Jabbari K et al. (2008). The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. *Nature* **456**: 239–244.
- Breton S, Jouhet J, Guyet U, Gros V, Pittera J et al. (2020). Unveiling membrane thermoregulation strategies in marine picocyanobacteria. *New Phytologist* **225**: 2396–2410.
- Brett MT, Müller-Navarra DC (1997). The role of highly unsaturated fatty acids in aquatic food web processes. *Freshwater Biology* **38**: 483–499.
- Brett MT, Müller-Navarra DC, Ballantyne A, Ravet JL, Goldman CR (2006). *Daphnia* fatty acid composition reflects that of their diet. *Limnology and Oceanography* **51**: 2428–2437.
- Budge SM, Iverson SJ, Bowen D, Ackman RG (2002). Among- and within-species variability in fatty acid signatures of marine fish and invertebrates on the Scotian Shelf, Georges Bank, and southern Gulf of St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences* **59**: 886–898.
- Calder PC (2015). Functional roles of fatty acids and their effects on human health. *Journal of Parenteral and Enteral Nutrition* **39**: 18S–32S.
- Chen CT, Liu Z, Ouellet M, Calon F, Bazinet RP (2009). Rapid β -oxidation of eicosapentaenoic acid in mouse brain: An in

- situ study. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* **80**: 157–163.
- Chen F, Johns MR (1991). Effect of C/N ratio and aeration on the fatty acid composition of heterotrophic *Chlorella sorokiniana*. *Journal of Applied Phycology* **3**: 203–209.
- Chorus I, Bartram J (1999). *Toxic Cyanobacteria in Water: A Guide to Their Public Health Consequences, Monitoring and Management*. London: World Health Organization, E&FN Spon, Routledge.
- Cohen Z, Vonshak A, Richmond A (1988). Effect of environmental conditions on fatty acid composition of the red alga *Porphyridium cruentum*: Correlation to growth. *Journal of Phycology* **2**: 328–332.
- Cook HW, McMaster CR (2004). Fatty acid desaturation and chain elongation in eukaryotes. In: Vance DE, Vance JE (eds.), *Biochemistry of Lipids, Lipoproteins and Membranes. New Comprehensive Biochemistry*, pp. 181–204. Amsterdam: Elsevier.
- Copeman LA, Parrish CC, Harel M, Brown JA (2002). Effects of docosahexaenoic, eicosapentaenoic, and arachidonic acids on the early growth, survival, lipid composition and pigmentation of yellowtail flounder (*Limanda ferruginea*): A live food enrichment experiment. *Aquaculture* **210**: 285–304.
- Dalsgaard J, St. John M, Kattner G, Mueller-Navarra DC, Hagen W (2003). Fatty acid trophic markers in the pelagic marine food environment. *Advances in Marine Biology* **46**: 225–234.
- de Castro Araujo S, Garcia VMT (2005). Growth and biochemical composition of the diatom *Chaetoceros* cf. *wighamii* brightwell under different temperature, salinity and carbon dioxide levels. I. Protein, carbohydrate and lipids. *Aquaculture* **246**: 405–412.
- de la Rosa F, de Troch M, Malanga G, Hernando M (2020). Differential sensitivity of fatty acids and lipid damage in *Microcystis aeruginosa* (cyanobacteria) exposed to increased temperature. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **235**: 108773.
- de Troch M, Boeckx P, Cnudde C, van Gansbeke D, Vanreusel A, Vincx M, Caramujo M (2012) Bioconversion of fatty acids at the basis of marine food webs: Insights from a compound specific stable isotope analysis. *Marine Ecology Progress Series* **465**: 53–67.
- Diau GY, Hsieh AT, Sarkadi-Nagy E, Wijendran V, Nathanielsz PW, Brenna JT (2005). The influence of long chain polyunsaturate supplementation on docosahexaenoic acid and arachidonic acid in baboon neonate central nervous system. *BMC Medicine* **3**: 11.
- Dodson VJ, Mouget JL, Dahmen JL, Leblond JD (2014). The long and short of it: Temperature-dependent modifications of fatty acid chain length and unsaturation in the galactolipid profiles of the diatoms *Haslea ostrearia* and *Phaeodactylum tricornerutum*. *Hydrobiologia* **727**: 95–107.
- Doney S, Ruckelshaus M, Duffy J, Barry J, Chan F et al. (2012). Climate change impacts on marine ecosystems. *Annual Review of Marine Science* **4**: 11–37.
- Drouin G, Rioux V, Legrand P (2019). The n-3 docosapentaenoic acid (DPA): A new player in the n-3 long chain polyunsaturated fatty acid family. *Biochimie* **159**: 36–48.
- Duarte AG, Katata G, Hoshika Y, Hossain M, Kreuzwieser J, Arneth A, Ruehr NK (2016). Immediate and potential long-term effects of consecutive heat waves on the photosynthetic performance and water balance in Douglas-fir. *Journal of Plant Physiology* **205**: 57–66. DOI 10.1016/j.jplph.2016.08.012.
- Ducklow HW, Fraser WR, Meredith MP, Stammerjohn S, Doney SC et al. (2013). West Antarctic Peninsula: An ice-dependent coastal marine ecosystem in transition. *Oceanography* **26**: 190–203. DOI 10.5670/oceanog.2013.62.
- Dunstan GA, Volkman JK, Barrett SM, Leroi JM, Jeffrey SW (1994). Essential polyunsaturated fatty acids from 14 species of diatom (Bacillariophyceae). *Phytochemistry* **35**: 155–161. DOI 10.1016/S0031-9422(00)90525-9.
- Dvoretzky AG, Bichkaeva FA, Baranova NF, Dvoretzky VG (2021). Fatty acid composition of the Barents Sea red king crab (*Paralithodes camtschaticus*) leg meat. *Journal of Food Composition and Analysis* **98**: 103826. DOI 10.1016/j.jfca.2021.103826.
- Economist (2013). <http://www.economist.com/news/science-and-technology/21571386>.
- Falkowski P, Barber R, Smetacek V (1998). Biogeochemical controls and feedbacks on ocean primary productivity. *Science* **281**: 200–206. DOI 10.1126/science.281.5374.200.
- Feijão E, Franzitta M, Cabrita MT, Caçador I, Duarte B, Gameiro C, Matos AR (2020). Marine heat waves alter gene expression of key enzymes of membrane and storage lipids metabolism in *Phaeodactylum tricornerutum*. *Plant Physiology and Biochemistry* **156**: 357–368.
- Feijão E, Gameiro C, Franzitta M, Duarte B, Caçador I, Cabrita MT, Matos AR (2017). Heat wave impacts on the model diatom *Phaeodactylum tricornerutum*: Searching for photochemical and fatty acid biomarkers of thermal stress. *Ecological Indicators* **95**: 1026–1037.
- Field CB, Behrenfeld MJ, Randerson JT, Falkowski P (1998). Primary production of the biosphere: Integrating terrestrial and oceanic components. *Science* **281**: 237–240.
- Flombaum P, Gallegos JL, Gordillo RA, Rincón J, Zabala LL et al. (2013). Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proceedings of the National Academy of Sciences, USA*, **110**: 9824–9829.
- Frölicher TL, Laufkötter C (2018). Emerging risks from marine heat waves. *Nature Communications* **9**: 2015–2018.
- Fuschino JR, Guschina IA, Dobson G, Yan ND, Harwood JL, Arts MT (2011). Rising water temperatures alter lipid dynamics and reduce n-3 essential FA concentrations in *Scenedesmus obliquus*. *Journal of Phycology* **47**: 763–774.
- Galloway AWE, Winder M (2015). Partitioning the relative importance of phylogeny and environmental conditions on phytoplankton fatty acids. *PLoS One* **10**: e0130053. DOI 10.1371/journal.pone.0130053.
- Gao K, Helbling EW, Häder DP, Hutchins DA (2012). Responses of marine primary producers to interactions between ocean acidification, solar radiation, and warming. *Marine Ecology Progress Series* **470**: 167–189. DOI 10.3354/meps10043.
- Gladyshev M, Arts MT, Sushchik N (2009). Preliminary estimates of the export of omega-3 highly unsaturated fatty acids (EPA + DHA) from aquatic to terrestrial ecosystems. In: Arts MT, Brett MT, Kainz MJ (eds.), *Lipids in Aquatic Ecosystems*, pp. 179–210. New York: Springer.
- Gladyshev M, Sushchik O, Makhutova N (2013). Production of EPA and DHA in aquatic ecosystems and their transfer to the land. *Prostaglandins Other Lipid Mediators* **107**: 117–126. DOI 10.1016/j.prostaglandins.2013.03.002.
- Gonçalves A, Marques JC, Gonçalves F (2017). Fatty acids' profiles of aquatic organisms: Revealing the impacts of environmental and anthropogenic stressors. *Fatty Acids* 89–118.
- Gurr MI (1985) Biosynthesis of fats. In: Padley FB, Podmore J (Eds.), *The Role of Fats in Human Nutrition*, pp. 24–28. Ellis Horwood, Ltd., Chichester.

- Guschina IA, Harwood JL (2009). Algal lipids and effect of the environment on their biochemistry. In: Arts MT, Brett MT, Kainz MJ (eds.), *Lipids in Aquatic Ecosystems*, pp. 1–24. New York: Springer.
- Häder DP, Gao K (2015). Interactions of anthropogenic stress factors on marine phytoplankton. *Frontiers in Environmental Science* **3**: 14.
- Häder DP, Williamson CE, Wangberg SA, Rautio M, Rose KC, Gao K, Helbling EW, Sinha RP, Worrest RP (2015). Effects of UV radiation on aquatic ecosystems and interactions with other environmental factors. *Photochemical & Photobiological Sciences* **14**: 108–126. DOI 10.1039/C4PP90035A.
- Hamilton M, Haslam R, Napier J, Sayanova O (2014). Metabolic engineering of microalgae for enhanced production of omega-3 long chain polyunsaturated fatty acids. *Metabolic Engineering* **22**: 3–9. DOI 10.1016/j.ymben.2013.12.003.
- Harwood JL (1998). Membrane lipids in algae. In: Siegenthaler PA, Murata N (eds.), *Lipids in Photosynthesis: Structure, Function and Genetics*, pp. 53–63. Dordrecht: Kluwer Academic Publishers.
- Harwood JL, Jones AL (1989). Lipid-metabolism in algae. *Advances in Botanical Research* **16**: 1–53. DOI 10.1016/S0065-2296(08)60238-4.
- Hazel JR (1995). Thermal adaptation in biological membranes: Is homeoviscous adaptation the explanation? *Annual Review of Physiology* **57**: 19–42. DOI 10.1146/annurev.ph.57.030195.000315.
- Henderson RJ, Hegseth EN, Park MT (1998). Seasonal variation in lipid and fatty acid composition of ice algae from the Barents Sea. *Polar Biology* **20**: 48–55. DOI 10.1007/s003000050275.
- Henley SF, Schofield OM, Hendry KR, Schloss IR, Steinberg DK et al. (2019). Variability and change in the west Antarctic Peninsula marine system: Research priorities and opportunities. *Progress in Oceanography* **173**: 208–237. DOI 10.1016/j.pocean.2019.03.003.
- Hernando M, Schloss IR, Almandoz GO, Malanga G, Varela DE, de Troch M (2018). Combined effects of temperature and salinity on fatty acid content and lipid damage in Antarctic phytoplankton. *Journal of Experimental Marine Biology and Ecology* **503**: 120–128. DOI 10.1016/j.jembe.2018.03.004.
- Hernando M, Schloss IR, Malanga G, Almandoz GO, Ferreyra GA, Aguiar MB, Puntarulo S (2015). Effects of salinity changes on coastal Antarctic phytoplankton physiology and assemblage composition. *Journal Experimental Marine Biology Ecology* **466**: 110–119. DOI 10.1016/j.jembe.2015.02.012.
- Hernando M, Varela DE, Malanga G, Almandoz GO, Schloss IR (2020). Effects of climate-induced changes in temperature and salinity on phytoplankton physiology and stress responses in coastal Antarctica. *Journal of Experimental Marine Biology and Ecology* **530**: 151400.
- Hixson SM, Sharma B, Kainz MJ, Wacker A, Arts MT (2015). Production, distribution, and abundance of long-chain omega-3 polyunsaturated fatty acids: A fundamental dichotomy between freshwater and terrestrial ecosystems. *Environmental Reviews* **23**: 414–424. DOI 10.1139/er-2015-0029.
- Hoffmann M, Marxen K, Schulz R, Vanselow KH (2010). TFA and EPA productivities of *Nannochloropsis salina* influenced by temperature and nitrate stimuli in turbidostatic controlled experiments. *Marine drugs* **8**: 2526–2545
- Huertas IE, Rouco M, Lopez-Rodas V, Costas E (2011). Warming will affect phytoplankton differently: Evidence through a mechanistic approach. *Proceedings of the Royal Society B: Biological Sciences* **278**: 3534–3543. DOI 10.1098/rspb.2011.0160.
- IPCC (2019). IPCC special report on the ocean and cryosphere in a changing climate. In: Pörtner HO, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al. Geneva: IPCC.
- Jiang H, Gao K (2004). Effects of lowering temperature during culture on the production of polyunsaturated fatty acids in the marine diatom *Phaeodactylum tricornutum* (Bacillariophyceae). *Journal of Phycology* **40**: 651–654.
- Jiang Y, Chen F (2000). Effects of temperature and temperature shift on docosahexaenoic acid production by the marine microalga *Cryptophyta ecodinium cohnii*. *Journal of the American Oil Chemists' Society* **77**: 613–617.
- Jónasdóttir S (2019). Fatty acid profiles and production in marine phytoplankton. *Marine Drugs* **17**: 151.
- Jónasdóttir SH, Visser AW, Jespersen C (2009). Assessing the role of food quality in the production and hatching of *Temora longicornis* eggs. *Marine Ecology Progress Series* **382**: 139–150.
- Kainz MJ, Arts MT, Mazumber A (2004). Essential fatty acids in the planktonic food web and their ecological role for higher trophic levels. *Limnology and Oceanography* **49**: 1784–1793.
- Kattner G, Hagen W (2009). Lipids in marine copepods: latitudinal characteristics and perspective to global warming. In: *Lipids in Aquatic Ecosystems*, pp. 257–280. Springer, New York, NY.
- Kaur G, Begg DP, Barr D, Garg M, Cameron-Smith D, Sinclair AJ (2010). Short-term docosapentaenoic acid (22: 5 n-3) supplementation increases tissue docosapentaenoic acid, DHA and EPA concentrations in rats. *British Journal of Nutrition* **103**: 32–37.
- Khazin-Goldberg I, Cohen Z (2006). The effect of phosphate starvation on the lipid and fatty acid composition of the fresh water eustigmatophyte *Monodus subterraneus*. *Phytochemistry* **67**: 696–701.
- Larsson U, Nyberg S, Zakrisson A, Hajdu S (2015). Baltic Sea phytoplankton: Long-term variability of major groups and primary production in spring and summer related to nutrients and temperature. <http://su.diva-portal.org/smash/record.jsf?pid=diva2%3A783358&dsid=-4719>.
- Li J, Han D, Wang D, Ning K, Jia J et al. (2014). Choreography of transcriptomes and lipidomes of *Nannochloropsis* reveals the mechanisms of oil synthesis in microalgae. *Plant Cell* **26**: 1645–1665.
- Liang Y, Beardall J, Heraud P (2006). Changes in growth, chlorophyll fluorescence and fatty acid composition with culture age in batch cultures of *Phaeodactylum tricornutum* and *Chaetoceros muelleri* (Bacillariophyceae). *Botanica Marina* **49**: 165–173.
- Lindsey R, Dahlman L (2017). Climate change: Global temperature. <https://www.climate.gov/news-features/understanding-climate/climate-change-globaltemperature>.
- Liu JJ, Green P, Mann JJ, Rapoport SI, Sublette ME (2015). Pathways of polyunsaturated fatty acid utilization: Implications for brain function in neuropsychiatric health and disease. *Brain Research* **1597**: 220–246.
- Los DA, Ray MK, Murata N (1997). Differences in the control of the temperature-dependent expression of four genes for desaturases in *Synechocystis* sp. PCC 6803. *Molecular Microbiology* **25**: 1167–1175.
- Maazouzi C, Masson G, Izquierdo MS, Pihan GC (2008). Midsummer heat wave effects on *Lacus trine* Plankton: Variation of assemblage structure and fatty acid composition. *Journal of Thermal Biology* **33**: 287–296.
- Maltse Y, Maltseva K (2021). Fatty acids of microalgae: Diversity and applications. *Reviews in Environmental Science and Bio-Technology* **20**: 515–547.

- Mansour MP, Frampton DMF, Nichols PD, Volkman JK, Blackburn SI (2005). Lipid and fatty acid yield of nine stationary-phase microalgae: Applications and unusual C-24-C-28 polyunsaturated fatty acids. *Journal of Applied Phycology* **17**: 287–300.
- Martin T (2000). Cyanobacterial dominance in lakes. *Hydrobiology* **438**: 1–12. DOI 10.1023/A:1004155810302.
- Massana R, Taylor LT, Murray AE, Wu KY, Jeffrey WH, DeLong EF (1998). Vertical distribution and temporal variation of marine planktonic archaea in the during early spring. *Limnology Oceanography* **43**: 607–617. DOI 10.4319/lo.1998.43.4.0607.
- Meredith MP, Stammerjohn SE, Venables HJ, Ducklow HW, Martinson DG et al. (2017). Changing distributions of sea ice melt and meteoric water west of the Antarctic Peninsula. *Deep Sea Research Part II: Topical Studies in Oceanography* **139**: 40–57. DOI 10.1016/j.dsr2.2016.04.019.
- Milledge JJ (2011). Commercial application of microalgae other than as biofuels: A brief review. *Reviews in Environmental Science and Biotechnology* **10**: 31–41. DOI 10.1007/s11157-010-9214-7.
- Mimouni V, Ulmann L, Pasquet V, Mathieu M, Picot L, Bougaran G, Cadoret JP, Morant-Manceau A, Schoefs B (2012). The potential of microalgae for the production of bioactive molecules of pharmaceutical interest. *Current Pharmaceutical Biotechnology* **13**: 2733–2750. DOI 10.2174/138920112804724828.
- Monroig Ó, Tocher DR, Navarro JC (2013). Biosynthesis of polyunsaturated fatty acids in marine invertebrates: Recent advances in molecular mechanisms. *Marine Drugs* **11**: 3998–4018.
- Morgan-Kiss RM, Ivanov AG, Modla S, Czymmek K, Hüner NPA, Priscu JC, Lisle JT, Hanson TE (2008). Identity and physiology of a new psychrophilic eukaryotic green alga, *Chlorella* sp., strain BI, isolated from a transitory pond near Bratina Island. *Antarctica Extremophiles* **12**: 701–711.
- Mortensen SH, Borsheim KY, Rainuzzo JK, Knutsen G (1988). Fatty acid and elemental composition of the marine diatom *Chaetoceros gracilis* Schutt. Effects of silicate deprivation, temperature and light intensity. *Journal Experimental Marine Biology and Ecology* **122**: 173–185.
- Mühlroth A, Li K, Røkke G, Winge P, Olsen Y, Hohmann-Marriott MF, Vadstein O, Bones AM (2013). Pathways of lipid metabolism in marine algae, co-expression network, bottlenecks and candidate genes for enhanced production of EPA and DHA in species of Chromista. *Marine Drugs* **11**: 4662–4697.
- Murray AE, Preston CM, Massana R, Taylor LT, Blakis A, Wu K, DeLong EF (1998). Seasonal and spatial variability of bacterial and archaeal assemblages in the coastal waters near Anvers Island. *Antarctica Applied and Environmental Microbiology* **64**: 2585–2595.
- Myhre G, Boucher O, Breon FM, Forster P, Shindell D (2015). Declining uncertainty in transient climate response as CO₂ forcing dominates future climate change. *Nature Geoscience* **8**: 181–185.
- Napolitano GE (1999). Fatty acids as trophic and chemical markers in freshwater ecosystems. In: Arts MT, Wainman BC (eds.), *Lipids in Freshwater Ecosystems*, pp. 21–44. New York: Springer.
- NOAA (2020). National Centers for Environmental Information, State of the Climate: Global Climate Report for Annual 2019. <https://www.ncdc.noaa.gov/sotc/global/201913>.
- Obata T, Fernie A, Nunes-Nesi A (2013). The central carbon and energy metabolism of marine diatoms. *Metabolites* **3**: 325–346.
- O'Donnell DR, Du ZY, Litchman E (2019). Experimental evolution of phytoplankton fatty acid thermal reaction norms. *Evolutionary applications* **12**: 1201–1211.
- Ohse S, Derner RB, Ozório RÁ, Corrêa RG, Furlong EB, Cunha PCR (2015). Lipid content and fatty acid profiles in ten species of microalgae. *Idesia* **33**: 93–101.
- Okuda T, Ando A, Negoro H, Muratsubaki T, Kikukawa H, Sakamoto T, Sakuradani E, Shimizu S, Ogawa J (2015). Eicosapentaenoic acid (EPA) production by an oleaginous fungus *Mortierella alpina* expressing heterologous the $\Delta 17$ -desaturase gene under ordinary temperature. *European Journal of Lipid Science and Technology* **117**: 1919–1927.
- Oliver RL, Gand GG (2000). Freshwater blooms. In: Whiton BA, Potts M (eds.), *The Ecology of Cyanobacteria*, pp. 149–194. Dordrecht: Kluwer Academic Press.
- Paerl HW, Huisman J (2008). Blooms like it hot. *Science* **320**: 57–58.
- Pal D, Khozin-Goldberg I, Cohen Z, Boussiba S (2011). The effect of light, salinity, and nitrogen availability on lipid production by *Nannochloropsis* sp. *Applied Microbiology Biotechnology* **90**: 1429–1441.
- Parrish C (2013). Lipids in marine ecosystems. *ISRN Oceanography* **2013**: 1–16. DOI 10.5402/2013/604045.
- Pasquet V, Ulmann L, Mimouni Guihéneuf F, Jacquette B, Manceau A, Gérard T (2014). Fatty acids profile and temperature in the cultured marine diatom *Odontella aurita*. *Journal of Applied Phycology* **26**: 2265–2271.
- Penfield S (2008). Temperature perception and signal transduction in plants. *New Phytologist* **179**: 615–628.
- Perkins-Kirkpatrick SE, Gibson PB (2017). Changes in regional heatwave characteristics as a function of increasing global temperature. *Scientific Reports* **7**: 12256.
- Poerschmann J, Spijkerman E, Langer U (2004). Fatty acid patterns in *Chlamydomonas* sp. as a marker for nutritional regimes and temperature under extremely acidic conditions. *Microbial Ecology* **48**: 78–89.
- Pussella PGRNI, Gunathilake J, Bandara KRMU, Dammalage T, Jayakody J (2015). Coastline changes: Vulnerability and predictions—A case study of the northwestern coastal belt of Sri Lanka. *Tourism, Leisure and Global Change* **2**: 125–154.
- Qiao H, Cong C, Sun C, Li B, Wang J, Zhang L (2016). Effect of culture conditions on growth, fatty acid composition and DHA/EPA ratio of *Phaeodactylum tricorutum*. *Aquaculture* **452**: 311–317.
- Raymont JE (2014). *Plankton & Productivity in the Oceans: Volume 1: Phytoplankton*. Elsevier, Pergamon Press, Oxford, England.
- Renaud SM, Parry DL (1994). Microalgae for use in tropical aquaculture II: Effect of salinity on growth, gross chemical composition and fatty acid composition of three species of marine microalgae. *Journal of Applied Phycology* **6**: 347–356.
- Renaud SM, Thinh LV, Lambrinidis G, Parry DL (2002). Effect of temperature on growth, chemical composition and fatty acid composition of tropical Australian microalgae grown in batch cultures. *Aquaculture* **211**: 195–214.
- Renaud SM, Zhou HC, Parry DL, Thinh LV, Woo KC (1995). Effect of temperature on the growth, total lipid content and fatty acid composition of recently isolated tropical microalgae *Isochrysis* sp., *Nitzschia closterium*, *Nitzschia paleacea*, and commercial species *Isochrysis* sp. (clone T.ISO). *Journal of Applied Phycology* **7**: 595–602.
- Reuss N, Poulsen LK (2002). Evaluation of fatty acids as biomarkers for a natural plankton community. A field study of a spring bloom and a post-bloom period off West Greenland. *Marine Biology* **141**: 423–434.

- Roncarati A, Meluzzi A, Acciarri S, Tallarico N, Meloti P (2004). Fatty acid composition of different microalgae strains (*Nannochloropsis* sp., *Nannochloropsis oculata* (Droop) Hibberd, *Nannochloris atomus* Butcher and *Isochrysis* sp.) according to the culture phase and the carbon dioxide concentration. *Journal of the World Aquaculture Society* **35**: 401–411.
- Rousch JM, Bingham SE, Sommerfeld MR (2003). Change in fatty acid profiles of thermos-intolerant and thermo-tolerant marine diatoms during temperature stress. *Journal of Experimental Marine Biology and Ecology* **295**: 145–156.
- Sakamoto T, Higashi S, Wada H, Murata N, Bryant DA (1997). Low-temperature-induced desaturation of fatty acids and expression of desaturase genes in the cyanobacterium *Synechococcus* sp. PCC 7002. *FEMS Microbiology Letters* **152**: 313–320.
- Sang M, Wang M, Liu J, Zhang C, Li A (2012). Effects of temperature, salinity, light intensity, and pH on the eicosapentaenoic acid production of *Pinguicoccus pyrenoidosus*. *Journal of Ocean University of China* **11**: 181–186.
- Sargent JR, Bell JG, Bell MV, Henderson RJ, Tocher DR (1993). The metabolism of phospholipids and polyunsaturated fatty acids in fish. In: Lahlou B, Vitiello P (eds.), *Aquaculture: Fundamental and Applied Research Coastal and Estuarine Studies*, pp. 103–112. Washington DC: American Geophysical Union.
- Schloss IR, Ferreyra GA, Ruiz-Pino D (2002). Phytoplankton biomass in antarctic shelf zones: A conceptual model based on Potter Cove, King George Island. *Journal of Marine Systems* **36**: 129–143.
- Shi H, Luo X, Wu R, Yue X (2018). Production of eicosapentaenoic acid by application of a delta-6 desaturase with the highest ALA catalytic activity in algae. *Microbial Cell Factories* **17**: 7.
- Shimajima M, Tsuchiya M, Ohta H (2009). Temperature-dependent hyper-activation of monoglucosyldiacylglycerol synthase is post-translationally regulated in *Synechocystis* sp. PCC 6803. *FEBS Letters* **583**: 2372–2376.
- Sinclair R (2000). Good, bad or essential fats: What is the story with Omega-3? *Nutrition & Food Science* **30**: 178–182.
- Sinensky M (1974). Homeoviscous adaptation—Homeostatic process that regulates viscosity of membrane lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* **71**: 522–525.
- Singh A, Ward OP (1997). Microbial production of docosahexaenoic acid (DHA, C22: 6). *Advances in Applied Microbiology* **45**: 271–312.
- Siron R, Giusti G, Berland B (1989). Changes in the fatty acid composition of *Phaeodactylum tricornerutum* and *Dunaliella tertiolecta* during growth and under phosphorus deficiency. *Marine Ecology Progress Series* **55**: 95–100.
- Springer M, Franke H, Pulz O (1994). Increase of the content of polyunsaturated fatty acids in *Porphyridium cruentum* by low-temperature stress and acetate supply. *Journal of plant physiology* **143**: 534–537.
- Steele NM, Fry SC (2000). Differences in catalytic properties between native isoenzymes of xyloglucanendotransglycosylase (XET). *Phytochemistry* **54**: 667–680.
- Stillwell W, Wassall SR (2003). Docosahexaenoic acid: Membrane properties of a unique fatty acid. *Chemistry and Physics of Lipids* **126**: 1–27.
- Stinson EE, Kwoczek R, Kurantz MJ (1991). Effect of cultural conditions on production of eicosapentaenoic acid by *Pythium irregulare*. *Journal of Industrial Microbiology* **8**: 171–178.
- Strasser RJ, Srivastava A, Tsimilli-Michael M (2000). The fluorescence transient as a tool to characterise and screen photosynthetic samples. In: Yunus M, Pathre U, Mohanty P (eds.), *Probing Photosynthesis: Mechanism, Regulation and Adaptation*, pp. 443–480. London: Taylor & Francis.
- Sukenik A (1991). Ecophysiological considerations in the optimization of eicosapentaenoic acid production by *Nannochloropsis* sp. (Eustigmatophyceae). *Bioresource Technology* **35**: 263–269.
- Sukenik A, Zamora O, Carmeli Y (1993). Biochemical quality of marine unicellular algae with emphasis on lipid composition. II. *Nannochloropsis* sp. *Aquaculture* **117**: 313–326.
- Sushchik NN, Kalacheva GS, Zhila NO, Gladyshev MI, Volova TG (2003). A temperature dependence of the intra- and extracellular fatty-acid composition of green algae and cyanobacterium. *Russian Journal of Plant Physiology (English Translation)* **50**: 374–380.
- Tassoni D, Kaur G, Weisinger RS, Sinclair AJ (2008). The role of eicosanoids in the brain. *Asia Pacific Journal of Clinical Nutrition* **17**: 220–228.
- Teoh ML, Chu WL, Marchant H, Phang SM (2004). Influence of culture temperature on the growth, biochemical composition and fatty acid profiles of six Antarctic microalgae. *Journal of Applied Phycology* **16**: 421–430. DOI 10.1007/s10811-004-5502-3.
- Teoh ML, Phang SM, Chu WL (2013). Response of Antarctic, temperate, and tropical microalgae to temperature stress. *Journal of Applied Phycology* **25**: 285–297. DOI 10.1007/s10811-012-9863-8.
- Thompson PA, Guo MX, Harrison PJ, Whyte JNC (1992). Effects of variation in temperature on the fatty-acid composition of 8 species of marine-phytoplankton. *Journal of Phycology* **28**: 488–497. DOI 10.1111/j.0022-3646.1992.00488.x.
- Thyssen M, Ferreyra G, Moreau S, Schloss I, Denis M, Demers S (2011). The combined effect of ultraviolet B radiation and temperature increase on phytoplankton dynamics and cell cycle using pulse shape recording flow cytometry. *Journal of Experimental Marine Biology and Ecology* **406**: 95–107. DOI 10.1016/j.jembe.2011.06.015.
- Tonon T, Harvey D, Larson TR, Graham IA (2002). Long chain polyunsaturated fatty acid production and partitioning to triacylglycerols in four microalgae. *Phytochemistry* **61**: 15–24. DOI 10.1016/S0031-9422(02)00201-7.
- Torstensson A, Jiménez C, Nilsson AK, Wulff A (2019). Elevated temperature and decreased salinity both affect the biochemical composition of the Antarctic sea-ice diatom *Nitzschia lecontei*, but not increased pCO₂. *Polar Biology* **42**: 2149–2164.
- Twining C, Bernhardt J, Derry A, Hudson CM, Ishikawa A et al. (2021). The evolutionary ecology of fatty-acid variation: Implications for consumer adaptation and diversification. *Ecology Letters* **24**: 1–23. DOI 10.1111/ele.13771.
- Twining CW, Brenna JT, Hairston NG, Flecker AS (2015). Highly unsaturated fatty acids in nature: What we know and what we need to learn. *Oikos* **125**: 749–760.
- Uauy R, Mena P, Rojas C (2000). Essential fatty acids in early life: Structural and functional role. *Proceedings of the Nutrition Society* **59**: 3–15.
- Vaezi R, Napier JA, Sayanova O (2013). Identification and functional characterization of genes encoding Omega-3 polyunsaturated fatty acid biosynthetic activities from unicellular microalgae. *Marine Drugs* **11**: 5116–5129.
- Van Doorslaer W, Stocks R, Jeppesen E, de Meester L (2007). Adaptive microevolutionary responses to simulated global

- warming in *Simocephalus vetulus*: A mesocosm study. *Global Change Biology* **13**: 878–886.
- Varkey D, Mazard S, Ostrowski M, Tetu SG, Haynes P, Paulsen IT (2016). Effects of low temperature on tropical and temperate isolates of marine *Synechococcus*. *ISME Journal* **10**: 1252–1263.
- Vincent WF (2000). Cyanobacterial dominance in the polar regions. In: *The Ecology of Cyanobacteria*, pp. 321–340. Dordrecht: Springer.
- Von Elert E (2004). Food quality constrains in *Daphnia*: Interspecific differences in the response to the absence of a long chain polyunsaturated fatty acid in the food source. *Hydrobiologia* **526**: 187–196.
- Wada H, Gombos Z, Murata N (1990). Enhancement of chilling tolerance of a cyanobacterium by genetic manipulation of fatty acid desaturation. *Nature* **347**: 200–203.
- Wada H, Murata N (1990). Temperature-induced changes in the fatty acids composition of the cyanobacterium, *Synechocystis* PCC 6803. *Plant Physiology* **92**: 1062–1069.
- Wen ZY, Chen F (2001). Application of statistically-based experimental designs for the optimization of eicosapentaenoic acid production by the diatom *Nitzschia leavis*. *Biotechnology and Bioengineering* **75**: 159–169.
- Winder M, Sommer U (2012). Phytoplankton response to a changing climate. *Hydrobiologia* **698**: 5–16.
- Winter R, Dzwolak W (2005). Exploring the temperature pressure configurational landscape of biomolecules: From lipid membranes to proteins. *Philosophical Transactions of the Royal Society of London Part A* **363**: 537–563.
- Xu XQ, Beardall J (1997). Effect of salinity on fatty acid composition of a green microalga from an antarctic hypersaline lake. *Phytochemistry* **45**: 655–658.
- Yang YH, Du L, Hosokawa M, Miyashita K, Kokubun Y, Arai H, Taroda H (2017). Fatty acid and lipid class composition of the microalga *Phaeodactylum tricorutum*. *Journal of Oleo Science* **66**: 363–368.
- Zainal Abidin ZA, Zainuddin Z, Wan Mastrai SFQ, Mohd Sidik Merican FM, Convey P (2020). Fatty acid profiles of Antarctic cyanobacteria *Leptolyngbya*. *Journal of Environmental Biology* **41**: 687–694.
- Zhu CJ, Lee YK, Chao TM (1997). Effects of temperature and growth phase on lipid and biochemical composition of *Isochrysis galbana* TK1. *Journal of Applied Phycology* **9**: 451–457. DOI 10.1023/A:1007973319348.
- Zulu NN, Zienkiewicz K, Vollheyde K, Feussner I (2018). Current trends to comprehend lipid metabolism in diatoms. *Progress in Lipid Research* **70**: 1–16. DOI 10.1016/j.plipres.2018.03.001.