

Truncated Bid Overexpression Induced by Recombinant Adenovirus Cre/LoxP System Suppresses the Tumorigenic Potential of CD133⁺ Ovarian Cancer Stem Cells

Qifang Long,* Weipei Zhu,* Jundong Zhou,† Jinchang Wu,† Weixian Lu,‡ Cui Zheng,‡
Dongmei Zhou,‡ Ling Yu,‡ and Ru Yang‡

*Department of Gynecology and Obstetrics, The Second Affiliated Hospital of Soochow University, Suzhou, Jiangsu Province, P.R. China

†Department of Radio-Oncology, Nanjing Medical University Affiliated Suzhou Hospital, Suzhou, Jiangsu Province, P.R. China

‡Department of Gynecology and Obstetrics, Nanjing Medical University Affiliated Suzhou Hospital, Suzhou, Jiangsu Province, P.R. China

Ovarian cancer is one of the most lethal malignant gynecologic tumors with a high relapse rate worldwide. Cancer stem cells (CSCs) have been identified in ovarian cancer and other malignant tumors as a small population of cells that are capable of self-renewal and multidifferentiation. CD133⁺ ovarian CSCs have been reported to be more tumorigenic and more resistant to chemotherapeutic treatment. Thus, CD133 has emerged as one of the most promising therapeutic markers for ovarian cancer treatment. In the current study, we constructed a recombinant adenovirus Cre/loxP regulation system to selectively introduce truncated Bid (tBid) expression specifically targeting CD133⁺ in ovarian CSCs. The results demonstrated that the coinfection of Ad-CD133-Cre and Ad-CMV-LoxP-Neo-LoxP-tBid significantly increased tBid expression in CD133⁺ ovarian CSCs. Moreover, the tBid overexpression induced by a recombinant adenovirus Cre/loxP system dramatically inhibited cell proliferation and invasion, significantly elevated cell apoptosis, and activated the mitochondrial apoptosis pathway in CD133⁺ ovarian CSCs. Additionally, recombinant adenovirus Cre/loxP system-mediated tBid overexpression suppressed the tumorigenic potential of CD133⁺ ovarian CSCs in a xenograft mouse model. In conclusion, our study successfully constructed a recombinant adenovirus Cre/loxP system and induced tBid overexpression in CD133⁺ ovarian CSCs, providing a new therapeutic approach for ovarian cancer treatment.

Key words: Truncated Bid (tBid); Cre/loxP system; Ovarian cancer; CD133; Cancer stem cells (CSCs)

INTRODUCTION

Ovarian cancer, one of the most lethal malignant gynecologic tumors, ranks as the fifth most common cause of cancer-related deaths, with 6.6 new cases per 100,000 women occurring worldwide each year¹. Early-stage ovarian cancer can be successfully removed by surgery alone, but approximately 70% of patients are diagnosed with ovarian cancer at either stage III or IV². At the advanced stages of ovarian cancer, surgical resection followed by platinum-based chemotherapy has been considered to be the standard treatment. However, most patients ultimately develop recurrence after treatment, which causes insurmountable obstacles for improving the survival rate^{3,4}. Despite the advances in surgical method

and chemotherapy agents, durable cures have not significantly increased in ovarian cancer treatment. Thus, it is urgent to develop effective treatment strategies to improve disease survival.

Cancer stem cells (CSCs; also called initiating cancer cells), a small subpopulation of cancer cells that possess multidifferentiation and self-renewal capacities, have been reported to be essential in cancer progression⁵. Populations of CSCs have been shown to be more tumorigenic in multiple experimental systems, including increased tumorigenicity, invasion, metastasis, angiogenic stimulation, and resistance to therapy⁶⁻⁹. It has been reported that CSCs were defined based on differential expression of cell surface markers (such as

CD117, CD119, CD24, CD44, and CD133)^{10,11}, differential biochemical properties (such as ALDH1)¹², and side population¹³. Ovarian cancer, like many cancers, is driven by CSCs. Increasing studies have demonstrated the carcinogenic effect of the CD133⁺ isolated from ovarian cancer cell lines or primary tumors¹⁴. The self-renewal, multidifferentiation, and tumor initiation capacities of CD133⁺ defined these cells as ovarian CSCs¹⁵. Therefore, the elimination of CD133⁺ CSCs may be a potential and promising therapeutic strategy for preventing ovarian cancer.

It has been reported that radiation and conventional chemotherapy target mostly the CD133⁻ population, evading the CD133⁺ population¹⁶. Therefore, the surviving CD133⁺ cells caused tumor relapse and reform after traditional therapy. Bid, a member of the BH3-only proteins, is activated via caspase 8-mediated cleavage in response to ligation of the death receptor¹⁷. Truncated Bid (tBid), the active form of Bid, translocates to the mitochondria from the cytosol, causing the activation of the mitochondrial death pathway and then inducing cell apoptosis^{18,19}. Several studies have suggested tBid to be a promising molecular target for killing cancer cells^{20,21}. To deliver tBid into cancer treatment, the recombinant adenoviruses have been suggested. Our study first constructed an adenovirus-mediated tBid expression initiated by the CMV promoter. However, recombinant adenovirus-induced tBid overexpression killed packaging cells during the production of recombinant adenovirus, which made it difficult to obtain high titers of recombinant adenovirus. Therefore, it is necessary to conduct a conditional tBid expression system.

In the present study, we constructed two recombinant adenoviruses, Ad-CD133-Cre and Ad-CMV-LoxP-Neo-LoxP-tBid, to achieve and investigate tBid overexpression in CD133⁺ ovarian CSCs. We hypothesized that Cre/loxP system-mediated tBid overexpression could specifically function in and kill CD133⁺ ovarian CSCs, thus providing a promising and effective method for preventing the metastasis and recurrence of ovarian cancer.

MATERIALS AND METHODS

Cell Culture

Ovarian cancer cell line SKOV-3 (ATCC[®] HTB-77[™]) was obtained from the American Type Culture Collection (ATCC; Manassas, VA, USA). SKOV-3 cells were cultured in McCoy's 5A medium (Thermo Fisher Scientific, Rockville, MD, USA) containing 10% fetal bovine serum (FBS; Gibco, Rockville, MD, USA), 0.1 mg/ml streptomycin, and 100 U/ml penicillin (Sigma-Aldrich, St. Louis, MO, USA) in 5% CO₂ at 37°C. The CD133⁺ ovarian CSCs were isolated from SKOV-3 cells using magnetic bead sorting as described previously¹⁴. Briefly, SKOV-3 cells

were dissociated with 0.25% trypsin (Sigma-Aldrich) and resuspended with ice-cold phosphate-buffered saline (PBS). The single cells were incubated with anti-CD133 monoclonal antibody-labeled microbeads (Miltenyi Biotech, Auburn, CA, USA) for 20 min at 4°C in the dark. Then samples were washed with ice-cold PBS and sorted on a BD fluorescence-activated cell sorting (FACS) Aria system (BD Biosciences, San Jose, CA, USA). The sorted CD133⁺ cells were cultured in a serum-free medium supplemented with 20 ng/ml human recombinant epidermal growth factor and 10 ng/ml human recombinant basic fibroblast growth factor (Invitrogen, Carlsbad, CA, USA).

Flow Cytometric Analysis

The purity of sorted CD133⁺ was detected using flow cytometric analysis. Briefly, cells were dissociated with trypsin, washed with ice-cold PBS, and resuspended into single-cell suspensions. Then cells were stained and incubated with mouse anti-human phycoerythrin (PE)-labeled CD133 (Miltenyi Biotech) for 30 min at 4°C in the dark. After incubation, the cells were washed with PBS and analyzed by FACS.

Construction, Production, and Purification of Recombinant Adenovirus

The cDNA fragments of the CD133 promoter and Cre were amplified and inserted into adenovirus pShuttle plasmid (Stratagene, Santa Clara, CA, USA) to construct the Ad-CD133-Cre recombinant adenovirus. The cDNA fragments of LoxP-Neo-LoxP and tBid were inserted into pShuttle-CMV plasmid (Stratagene) to construct the Ad-CMV-LoxP-Neo-LoxP-tBid recombinant adenovirus. The recombinant pShuttle plasmids and pAdEasy-1 plasmids (Stratagene) were homologously recombined in BJ5183 bacteria. The recombined plasmids were linearized and transfected into 293T cells (ATCC) to produce the recombinant virus. After 14 days, cells were harvested and lysed by freeze-thawing. The supernatants were collected and concentrated by CsCl gradient centrifugation. The titers were determined using the 50% tissue culture infectious dose method²². CD133⁺ cells were infected with Ad-CD133-Cre at 20 multiplicity of infection (MOI) and Ad-CMV-LoxP-Neo-LoxP-tBid at 30 MOI.

Luciferase Reporter Assay

Sorted CD133⁺ cells were seeded in 24-well plates. Twenty-four hours later, indicated luciferase reporter plasmids plus 3 ng of pRL-TK *Renilla* plasmid were transfected into the cells using Lipofectamine 2000 reagent (Life Technologies, Carlsbad, CA, USA) for 48 h. The luciferase activity was detected by a fluorescence detector (Promega, Madison, WI, USA).

RNA Extraction and Quantitative Real-Time PCR (qRT-PCR)

Total RNA was extracted from sorted CD133⁺ cells using TRIzol reagent (Invitrogen). The concentration and purity of total RNA were determined using an ultraviolet spectrophotometer (Eppendorf, Hamburg, German). The expression of mRNA was analyzed by real-time PCR using primers synthesized by the Shanghai Sangon Biological Engineering and Technology Service (Shanghai, P.R. China). Briefly, 20 μ l of reactions containing 50 ng of total RNA, 10 μ l of 2 \times SYBR Green PCR Master Mix, 6.25 μ l of AMV reverse transcriptase, 10 μ l of RNase inhibitor, and 0.1 mM primers was subjected to one cycle of 95°C for 10 min and then 40 cycles of 95°C for 15 s, 56°C for 30 s, and 72°C for 45 s. The expression of mRNA was normalized to β -actin. Data were analyzed by the Δ Ct method and expressed as fold changes.

Western Blot

For cytochrome c detection, cytosolic and mitochondrial proteins were extracted using a mitochondria isolation kit (Pierce, Rockford, IL, USA). For other proteins, the total proteins from the sorted CD133⁺ cells were extracted using RIPA lysis buffer (Beyotime, Nantong, P.R. China) and quantified with the BCA assay (Pierce). The proteins were loaded and separated by 12.5% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred onto a nitrocellulose membrane (Bio-Rad, Hercules, CA, USA). After blocking in 2.5% nonfat milk at 37°C for 1 h, the membrane was blotted with primary mouse anti-human antibodies: anti-tBid, anti-cytochrome c, anti-Bax, anti-Bcl-2, anti-caspase 3, anti-caspase 9, and anti- β -actin (Santa Cruz Biotechnology, Santa Cruz, CA, USA) at 4°C overnight. The membranes were then incubated with horseradish peroxidase (HRP)-conjugated secondary antibody (1:2,000; Bioss Antibodies, Beijing, P.R. China) for 1 h at room temperature. A fluorescent Western blotting detection system was used. The band density of each gene was normalized to the corresponding density of β -actin.

MTT Assay

Cell viability was detected using a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. In brief, cells were plated onto a 96-well culture plate at a density of 1×10^4 cells/well and cultured overnight. Thereafter, cells were infected with Ad-CD133-Cre or/and Ad-CMV-LoxP-Neo-LoxP-tBid and incubated for 48 h. The medium was replaced with fresh medium, and 20 μ l of MTT stock solution (5 mg/ml; Sigma-Aldrich) was added to each well. After culturing for 4 h, 200 μ l of dimethyl sulfoxide was added to each well to dissolve

the formazan crystal. Then absorbance at a wavelength of 490 nm was detected using a microplate reader (Bio-Tek Instruments, Winooski, VT, USA).

Transwell Assay

The invasive behaviors of the indicated cells were analyzed by Transwell chamber (Corning Costar-Corp., Cambridge, MA, USA) assay. In brief, 2×10^5 transfected cells in serum-free medium were added to each upper chamber precoated with Matrigel matrix, and 500 μ l of McCoy's 5A medium containing 10% FBS was added to the lower chamber to serve as chemoattractant. After 24 h of incubation, cells on the surface of the upper chamber were removed by scraping with a cotton swab, and the invasive cells on the surface of the lower membrane were fixed with 4% paraformaldehyde, stained with 0.1% crystal violet, photographed, and quantified by counting them in five random fields by a light microscope (Olympus, Tokyo, Japan).

Apoptosis Analysis

The apoptotic ratios of cells were determined using Annexin-V-FITC/propidium iodide (PI) apoptosis detection kits (Invitrogen, Shanghai, P.R. China). In brief, cells were first dissociated with trypsin, washed with ice-cold PBS, and resuspended in binding buffer at 1×10^6 cells/ml. Then the cells were incubated with annexin V and PI stock solution at 4°C in the dark. Thereafter, the samples were detected by FACS, and the data were analyzed by FlowJo software (BD Biosciences).

Animals

Six-week-old female severe combined immunodeficient (SCID) mice, weighing 20–28 g, were purchased from Guangdong Medical Laboratory Animal Center (Guangzhou, P.R. China). All experimental animals were housed under specific pathogen-free conditions ($21 \pm 2^\circ\text{C}$ and 12 h/12 h) with free access to water and food. The animal experimental procedures were approved by the Institutional Animal Care and Use Committee of Nanjing Medical University Affiliated Suzhou Hospital.

In Vivo Xenograft Experiments

The CD133⁺ ovarian CSCs (1×10^5 cells) were suspended in 200 μ l of PBS and injected subcutaneously into the right flank of SCID mice ($n=6$ mice/group). About 1×10^{10} plaque-forming units of recombinant adenovirus diluted in 50 μ l of PBS were injected intratumorally every 3 days. Tumor volume was measured using an external caliper and calculated according to the formula: $\text{length} \times \text{width}^2 \times \pi/6$. The mice were sacrificed by subcutaneous injection with sodium pentobarbital (100 mg/kg) 30 days after inoculation, and the tumors were excised for Western blot analysis.

Statistical Analysis

All the experiments in this study were performed three times. All data from this study were obtained from at least three independent experiments and were expressed as mean \pm SD. All statistical analyses were carried out using the software SPSS version 11.0 for Windows (SPSS Inc., Chicago, IL, USA). Statistical analysis between two samples was performed using Student's *t*-test. The chi-square or Fisher's exact tests were used to evaluate the differences in proportion. Differences were considered significant with a value of $p < 0.05$.

RESULTS

Coinfection of Ad-CD133-Cre and Ad-CMV-Loxp-Neo-Loxp-tBid Elevates tBid Expression in CD133⁺ Ovarian Cancer Stem Cells

To investigate the effect of tBid overexpression in CD133⁺ ovarian CSCs, the CD133⁺ cells were sorted from the total SKOV-3 cell line. The purity of CD133⁺ was 99.97 in sorted cells and 0.94 in total cells, respectively (Fig. 1A). To achieve tBid overexpression in CD133⁺

ovarian CSCs, we introduced a Cre/loxP-mediated expression system. We first inserted the open reading frame of Cre downstream of the CD133 promoter, which was cloned into an adenovirus (Fig. 1B). Therefore, the Cre would only be expressed when the CD133 promoter was activated. Then we constructed another recombinant adenovirus containing a flanked loxP cassette upstream of tBid. In this way, tBid would only be expressed after deleting the loxP cassette. After a pretest, the CD133⁺ ovarian CSCs were infected with Ad-CD133-Cre and Ad-CMV-Loxp-Neo-Loxp-tBid at the proportion of 3:2. The transfection efficiency was measured by luciferase activity report after the adenoviruses were marked with green fluorescent protein genes. The results demonstrated that recombinant adenoviruses were successfully transfected into CD133⁺ cells (Fig. 1C and E). The effect of the Cre/loxP system, the mRNA (Fig. 1F), and protein expression (Fig. 1D and G) of tBid was detected. The results demonstrated that tBid expression levels were significantly increased in the Ad-CD133-Cre and Ad-CMV-Loxp-Neo-Loxp-tBid coinfection group compared with the control group. Additionally, the single infection of




Figure 1. Cotransfection of Ad-CD133-Cre and Ad-CMV-Loxp-Neo-Loxp-tBid elevates tBid expression in CD133⁺ ovarian cancer stem cells. (A) The percentage of CD133⁺ in total SKOV-3 cell line and sorted CD133⁺ were measured by flow cytometry. (B) Diagram of Cre/loxP system-mediated tBid overexpression. The recombinant adenoviruses were infected into CD133⁺ ovarian cancer stem cells: Control group, CD133⁺ cells; Empty Ad group, CD133⁺ cell infected with empty adenovirus; Ad-Cre group, CD133⁺ cell infected with Ad-CD133-Cre; Ad-Loxp-tBid group, CD133⁺ cell infected with Ad-CMV-Loxp-Neo-Loxp-tBid; Ad-Cre + Ad-Loxp-tBid group, CD133⁺ cell coinfecting with Ad-CD133-Cre and Ad-CMV-Loxp-Neo-Loxp-tBid. (C, E) The adenoviruses infection efficiency was detected by luciferase activity. (F) The mRNA expression level of tBid was measured by qRT-PCR and normalized to β -actin. (D, G) The protein expression level of tBid was measured by Western blot assay and normalized to β -actin. *** $p < 0.001$ versus Control group.

Ad-CD133-Cre or Ad-CMV-Loxp-Neo-LoxP-tBid did not result in tBid overexpression. These results illustrated that Bid overexpression was successfully induced by the recombinant adenovirus Cre/loxP system in CD133⁺ ovarian CSCs.

Recombinant Adenovirus Cre/LoxP System-Induced tBid Overexpression Inhibits Cell Proliferation and Invasion of CD133⁺ Ovarian Cancer Stem Cells

To evaluate the effect of recombinant adenovirus Cre/loxP system-induced tBid overexpression, the CD133⁺ ovarian CSCs were coinfecting with Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid. The number of cells was obviously decreased in the coinfecting group compared with the control or single recombinant adenovirus infection group (Fig. 2). The cell proliferation rate was measured by the MTT method, and results showed that a single recombinant adenovirus infection had no distinct effect on cell proliferation. Consistent with the former results, coinfection of Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid significantly inhibited the cell proliferation of CD133⁺ ovarian CSCs. The invasion ability of CD133⁺ ovarian CSCs was measured by the Transwell assay. The results demonstrated that cell invasion was significantly decreased in the Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid coinfection group compared with the other groups.

Recombinant Adenovirus Cre/LoxP System-Induced tBid Overexpression Increases Cell Apoptosis of CD133⁺ Ovarian CSCs

We further investigate the effect of Cre/loxP system-induced tBid overexpression on the apoptotic rate of CD133⁺ ovarian CSCs. Cell apoptosis was detected by flow cytometric analysis using annexin V-FITC/PI apoptosis detection kits (Fig. 3). The results showed that the coinfection of Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid dramatically increased the apoptotic rate in CD133⁺ ovarian CSCs.

Recombinant Adenovirus Cre/LoxP System-Induced tBid Overexpression Activates the Mitochondrial Apoptosis Pathway in CD133⁺ Ovarian Cancer Stem Cells

It is well known that tBid activated the mitochondrial apoptosis pathway and induced cell death in various cancers²³. To verify the effect of recombinant adenovirus Cre/loxP system-induced tBid overexpression, we assessed the expression of the mitochondrial apoptosis pathway. The results demonstrated that coinfection of Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid significantly upregulated the release of cytochrome c from the mitochondria into the cytosol compared with the other groups. Furthermore, the proapoptosis proteins Bcl-2, caspase 3,




Figure 2. Recombinant adenovirus Cre/loxP system-induced tBid overexpression inhibits cell proliferation and invasion of CD133⁺ ovarian cancer stem cells. (A) The cells infected with Ad-CD133-Cre and Ad-CMV-Loxp-Neo-LoxP-tBid and cultured in serum-free medium containing 10 ng/ml human recombinant basic fibroblast growth factor and 20 ng/ml human recombinant epidermal growth factor. (B) Cell proliferation was detected by MTT assay. (C, D) Cell invasion ability was measured by the Transwell assay. ****p* < 0.001 versus Control group.




Figure 3. Recombinant adenovirus Cre/loxP system-induced tBid overexpression increases cell apoptosis of CD133⁺ ovarian cancer stem cells. (A) Cell apoptosis was detected by annexin V-FITC/PI assay. (B) Quantitative statistics of cell apoptosis rate. *** $p < 0.001$ versus Control group.




Figure 4. Recombinant adenovirus Cre/loxP system-induced tBid overexpression activates mitochondrial apoptosis pathway in CD133⁺ ovarian cancer stem cells. (A) Protein expression level was measured by the Western blot method. (B) Relative protein expression was quantified using Image-Pro Plus 6.0 software and normalized to β -actin. * $p < 0.005$ versus Control group.

and caspase 9 were significantly increased, while the antiapoptosis protein Bax was obviously decreased in the coinfection group (Fig. 4).

Recombinant Adenovirus Cre/LoxP System-Induced tBid Overexpression Ameliorates Ovarian Cancer In Vivo

To investigate the effect of recombinant adenovirus Cre/loxP system-induced tBid overexpression on tumor growth in vivo, the CD133⁺ ovarian CSCs were coinfecting with Ad-CD133-Cre and Ad-CMV-LoxP-Neo-LoxP-tBid and injected into the right scapula of SCID mice. SCID mice injected with blank CD133⁺ ovarian CSCs were used as the control. The results showed that mice in the coinfection group exhibited a higher survival rate with significantly smaller tumor volume compared with the control group (Fig. 5A–C). Moreover, the tBid expression level was dramatically increased in the coinfection group (Fig. 5D). These results indicated that adenovirus Cre/loxP system-induced tBid

overexpression suppressed the CD133-induced cancer growth in vivo.

DISCUSSION

Since the discovery of ovarian CSCs, the identification of strategies aimed at eliminating CSCs is of great interest to improve the survival of ovarian cancer²⁴. In the current study, we successfully constructed a recombinant adenovirus Cre/loxP system and achieved conditional tBid overexpression specifically targeting CD133⁺ ovarian CSCs. Moreover, the tBid overexpression induced by the recombinant adenovirus Cre/loxP system significantly inhibited the tumorigenic potential of CD133⁺ ovarian CSCs in in vitro and in a xenograft model. Taken together, our results reveal a novel adenovirus Cre/loxP system to accomplish tBid overexpression and provide a promising therapeutic strategy in ovarian cancer treatment.

It has been reported that CSCs were associated with tumor initiation, metastasis, chemoresistance, and recurrence²⁵. CD133 has been suggested as a stem cell marker




Figure 5. Recombinant adenovirus Cre/loxP system-induced tBid overexpression ameliorates ovarian cancer in vivo. Control group, SCID mice injected with CD133⁺ ovarian cancer stem cells; Ad-Cre+Ad-Loxp-tBid group, SCID mice injected with CD133⁺ ovarian cancer stem cell coinfecting with Ad-CD133-Cre and Ad-CMV-LoxP-Neo-LoxP-tBid. (A) The survival percentage of SCID mice. (B) The tumor growth curve of SCID mice. (C) Top image of representative tumors from SCID mice treated with different CD133⁺ ovarian cancer stem cells. (D) The protein expression of tBid was detected by Western blot and normalized to β -actin. * $p < 0.005$ versus Control group.

for various cancers^{26–29}. Expression of CD133 in cancer-initiating cells is well documented for prostate, brain, breast, and colon cancers, and is indicative of a poor prognosis in many malignant tumors. Ovarian cancer cell lines and primary tumors have been characterized for the expression of several CSC biochemical markers, in which CD133 emerged as the most promising³⁰. In the current study, CD133⁺ CSCs were successfully isolated from the SKOV-3 ovarian cancer cell line, and the purity of CD133⁺ reached 99.97 in sorted cells. Studies have shown that CD133⁺ ovarian CSCs are more tumorigenic and more resistant to chemotherapeutic treatment¹⁴. Thus, CD133⁺ CSCs were an effective and promising therapeutic target for ovarian cancer treatment. However, the method for eliminating CD133⁺ CSCs was limited due to the complex molecular structure of CD133.

In this study, we constructed a Cre/loxP regulation system to selectively augment tBid expression, specifically targeting CD133⁺ ovarian CSCs. It has been reported that the carcinoembryonic antigen (CEA) promoter-mediated Cre/loxP system induces the expression of the cytosine deaminase and specifically inhibits CEA-producing gastric cancer cells *in vitro* and *in vivo*³¹. Similarly, we used a CD133 promoter to control the expression of selective genes. In the current study, we wanted to introduce the suicide gene tBid into CD133⁺ ovarian CSCs and achieve the goal of eliminating CD133⁺ cells. However, it is hard to obtain a recombinant adenovirus carrying tBid driven by the CMV potent promoter, due to the cell killing caused by tBid expression during the recombinant adenovirus production. Thus, we needed to construct a conditional tBid expression system. Therefore, the Cre/loxP regulation system was used to elevate the selective expression of tBid. Cre is a site-specific recombinase that can mediate the specific recombinant process between two loxP sequences^{32,33}. We first constructed Ad-CMV-LoxP-Neo-LoxP-tBid, a recombinant adenovirus carrying tBid (with CMV promoter), and a null gene expression cassette (Neo) flanked by loxP was inserted upstream of tBid to block tBid expression. To conditionally initiate tBid expression in CD133⁺ stem cells, we constructed another recombinant adenovirus, Ad-CD133-Cre, carrying Cre with a CD133 promoter inserted upstream of Cre. In this way, the infection of Ad-CD133-Cre into CD133⁺ ovarian CSCs would trigger the expression of Cre, which would then mediate the loxP recombinant and cut off the null gene expression cassette between the two loxP, ultimately leading to tBid overexpression³⁴. In our study, coinfection of Ad-CMV-LoxP-Neo-LoxP-tBid and Ad-CD133-Cre significantly elevated the expression of tBid, suggesting that we successfully introduced tBid into CD133⁺ ovarian CSCs.

tBid is normally reported to induce cell apoptosis in various cancer both *in vivo* and *in vitro*³⁵. The results

in our study demonstrated that tBid overexpression was induced by the recombinant adenovirus Cre/loxP system, and it not only significantly increased cell apoptosis but also inhibited cell proliferation and invasion. This suggested that Cre/loxP system-mediated tBid overexpression suppressed the tumorigenic potential of CD133⁺ ovarian CSCs. Moreover, the coinfection of Ad-CMV-LoxP-Neo-LoxP-tBid and Ad-CD133-Cre dramatically increased the mitochondrial release of cytochrome c; promoted the expression of proapoptotic protein Bax, caspase 3, and caspase 9; and inhibited the antiapoptotic protein Bcl-2. These results suggested that Cre/loxP system-mediated tBid overexpression activated the mitochondrial apoptosis pathway in CD133⁺ ovarian CSCs. Additionally, we verified the effect of adenovirus Cre/loxP system-mediated tBid overexpression in a xenograft model. The results were consistent with the *in vivo* study, further identifying that Cre/loxP system-mediated tBid overexpression increased the survival of CD133⁺-induced ovarian cancer.

In summary, the present study established and applied the adenovirus Cre/loxP system to introduce selective tBid expression in CD133⁺ ovarian CSCs. The overexpression induced by the adenovirus Cre/loxP system significantly inhibited the tumorigenic potential of CD133⁺ ovarian CSCs both *in vivo* and in the xenograft model, providing a new therapeutic approach for ovarian cancer treatment.

ACKNOWLEDGMENTS: This study was supported by grants from Natural Science Foundation of Jiangsu Province (BK2012599), Suzhou Municipal Science and Technology Development Project (SYS2012088), and Suzhou Municipal Science and Education and Health Project (SZZX201506). Our study was approved by the ethics committee of the Institutional Animal Care and Use Committee of Nanjing Medical University Affiliated Suzhou Hospital. The authors declare no conflicts of interest.

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin.* 2016;66:7–30.
2. Yang D, Sun Y, Hu L, Zheng H, Ji P, Pecot CV, Zhao Y, Reynolds S, Cheng H, Rupaimoole R, Cogdell D, Nykter M, Broaddus R, Rodriguez-Aguayo C, Lopez-Berestein G, Liu J, Shmulevich I, Sood AK, Chen K, Zhang W. Integrated analyses identify a master microRNA regulatory network for the mesenchymal subtype in serous ovarian cancer. *Cancer Cell* 2013;23:186–99.
3. Chekerov R, Braicu I, Castillo-Tong DC, Richter R, Cadron I, Mahner S, Woelber L, Marth C, Van Gorp T, Speiser P, Zeillinger R, Vergote I, Sehouli J. Outcome and clinical management of 275 patients with advanced ovarian cancer International Federation of Obstetrics and Gynecology II to IV inside the European Ovarian Cancer Translational Research Consortium-OVCAD. *Int J Gynecol Cancer* 2013; 23:268–75.
4. Peiretti M, Bristow RE, Zapardiel I, Gerardi M, Zanagnolo V, Biffi R, Landoni F, Bocciolone L, Aletti GD, Maggioni A. Rectosigmoid resection at the time of primary cytoreduction

- for advanced ovarian cancer. A multi-center analysis of surgical and oncological outcomes. *Gynecol Oncol.* 2012; 126:220–3.
5. Szeke B, Sillo P, Fabian M, Mayer B, Karpati S, Nemeth K. Role of cancer stem cells in the progression and heterogeneity of melanoma. *Orv Hetil.* 2016;157:1339–48.
 6. Singh SK, Clarke ID, Terasaki M, Bonn VE, Hawkins C, Squire J, Dirks PB. Identification of a cancer stem cell in human brain tumors. *Cancer Res.* 2003;63:5821–8.
 7. Zhang S, Balch C, Chan MW, Lai HC, Matei D, Schilder JM, Yan PS, Huang TH, Nephew KP. Identification and characterization of ovarian cancer-initiating cells from primary human tumors. *Cancer Res.* 2008;68:4311–20.
 8. Iwasaki H, Suda T. Cancer stem cells and their niche. *Cancer Sci.* 2009;100:1166–72.
 9. Zhao J. Cancer stem cells and chemoresistance: The smartest survives the raid. *Pharmacol Ther.* 2016;160:145–58.
 10. Gao MQ, Choi YP, Kang S, Youn JH, Cho NH. CD24⁺ cells from hierarchically organized ovarian cancer are enriched in cancer stem cells. *Oncogene* 2010;29:2672–80.
 11. Curley MD, Therrien VA, Cummings CL, Sergent PA, Koulouris CR, Friel AM, Roberts DJ, Seiden MV, Scadden DT, Rueda BR, Foster R. CD133 expression defines a tumor initiating cell population in primary human ovarian cancer. *Stem Cells* 2009;27:2875–83.
 12. Silva IA, Bai S, McLean K, Yang K, Griffith K, Thomas D, Ginestier C, Johnston C, Kueck A, Reynolds RK, Wicha MS, Buckanovich RJ. Aldehyde dehydrogenase in combination with CD133 defines angiogenic ovarian cancer stem cells that portend poor patient survival. *Cancer Res.* 2011;71:3991–4001.
 13. Moserle L, Indraccolo S, Ghisi M, Frasson C, Fortunato E, Canevari S, Miotti S, Tosello V, Zamarchi R, Corradin A, Minuzzo S, Rossi E, Basso G, Amadori A. The side population of ovarian cancer cells is a primary target of IFN- α antitumor effects. *Cancer Res.* 2008;68:5658–68.
 14. Long H, Xie R, Xiang T, Zhao Z, Lin S, Liang Z, Chen Z, Zhu B. Autocrine CCL5 signaling promotes invasion and migration of CD133⁺ ovarian cancer stem-like cells via NF- κ B-mediated MMP-9 upregulation. *Stem Cells* 2012;30:2309–19.
 15. Baba T, Convery PA, Matsumura N, Whitaker RS, Kondoh E, Perry T, Huang Z, Bentley RC, Mori S, Fujii S, Marks JR, Berchuck A, Murphy SK. Epigenetic regulation of CD133 and tumorigenicity of CD133⁺ ovarian cancer cells. *Oncogene* 2009;28:209–18.
 16. Hossain M, Banik NL, Ray SK. Synergistic anti-cancer mechanisms of curcumin and paclitaxel for growth inhibition of human brain tumor stem cells and LN18 and U138MG cells. *Neurochem Int.* 2012;61:1102–13.
 17. Choi ES, Chung T, Kim JS, Lee H, Kwon KH, Cho NP, Cho SD. Mithramycin A induces apoptosis by regulating the mTOR/Mcl-1/tBid pathway in androgen-independent prostate cancer cells. *J Clin Biochem Nutr.* 2013;53:89–93.
 18. Esposti MD. The roles of Bid. *Apoptosis* 2002;7:433–40.
 19. Luo X, Budihardjo I, Zou H, Slaughter C, Wang X. Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* 1998;94:481–90.
 20. Hu BG, Liu LP, Chen GG, Ye CG, Leung KK, Ho RL, Lin MC, Lai PB. Therapeutic efficacy of improved alpha-fetoprotein promoter-mediated tBid delivered by folate-PEI600-cyclodextrin nanopolymer vector in hepatocellular carcinoma. *Exp Cell Res.* 2014;324:183–91.
 21. Cristofanon S, Fulda S. ABT-737 promotes tBid mitochondrial accumulation to enhance TRAIL-induced apoptosis in glioblastoma cells. *Cell Death Dis.* 2012;3:e432.
 22. Kanegae Y, Makimura M, Saito I. A simple and efficient method for purification of infectious recombinant adenovirus. *Jpn J Med Sci Biol.* 1994;47:157–66.
 23. Tan CT, Zhou QL, Su YC, Fu NY, Chang HC, Tao RN, Sukumaran SK, Baksh S, Tan YJ, Sabapathy K, Yu CD, Yu VC. MOAP-1 mediates Fas-induced apoptosis in liver by facilitating tBid recruitment to mitochondria. *Cell Rep.* 2016;16:174–85.
 24. Skubitz AP, Taras EP, Boylan KL, Waldron NN, Oh S, Panoskaltzis-Mortari A, Vallera DA. Targeting CD133 in an in vivo ovarian cancer model reduces ovarian cancer progression. *Gynecol Oncol.* 2013;130:579–87.
 25. Zhang QH, Dou HT, Xu P, Zhuang SC, Liu PS. Tumor recurrence and drug resistance properties of side population cells in high grade ovary cancer. *Drug Res. (Stuttg)* 2015;65:153–7.
 26. Peichev M, Naiyer AJ, Pereira D, Zhu Z, Lane WJ, Williams M, Oz MC, Hicklin DJ, Witte L, Moore MA, Rafii S. Expression of VEGFR-2 and AC133 by circulating human CD34(+) cells identifies a population of functional endothelial precursors. *Blood* 2000;95:952–8.
 27. Uchida N, Buck DW, He D, Reitsma MJ, Masek M, Phan TV, Tsukamoto AS, Gage FH, Weissman IL. Direct isolation of human central nervous system stem cells. *Proc Natl Acad Sci USA* 2000;97:14720–5.
 28. Corbeil D, Roper K, Hellwig A, Tavian M, Miraglia S, Watt SM, Simmons PJ, Peault B, Buck DW, Huttner WB. The human AC133 hematopoietic stem cell antigen is also expressed in epithelial cells and targeted to plasma membrane protrusions. *J Biol Chem.* 2000;275:5512–20.
 29. Weigmann A, Corbeil D, Hellwig A, Huttner WB. Prominin, a novel microvilli-specific polytopic membrane protein of the apical surface of epithelial cells, is targeted to plasmalemmal protrusions of non-epithelial cells. *Proc Natl Acad Sci USA* 1997;94:12425–30.
 30. Liang J, Yang B, Cao Q, Wu X. Association of vasculogenic mimicry formation and CD133 expression with poor prognosis in ovarian cancer. *Gynecol Obstet Invest.* 2016; 81(6):529–36.
 31. Ueda K, Iwahashi M, Nakamori M, Nakamura M, Matsuura I, Yamaue H, Tanimura H. Carcinoembryonic antigen-specific suicide gene therapy of cytosine deaminase/5-fluorocytosine enhanced by the cre/loxP system in the orthotopic gastric carcinoma model. *Cancer Res.* 2001; 61:6158–62.
 32. Stricklett PK, Nelson RD, Kohan DE. Site-specific recombination using an epitope tagged bacteriophage P1 Cre recombinase. *Gene* 1998;215:415–23.
 33. Kanegae Y, Lee G, Sato Y, Tanaka M, Nakai M, Sakaki T, Sugano S, Saito I. Efficient gene activation in mammalian cells by using recombinant adenovirus expressing site-specific Cre recombinase. *Nucleic Acids Res.* 1995; 23:3816–21.
 34. Zhou L, Wei X, Cheng L, Tian J, Jiang JJ. CD133, one of the markers of cancer stem cells in Hep-2 cell line. *Laryngoscope* 2007;117:455–60.
 35. Chaurio RA, Munoz LE, Maueroeder C, Janko C, Harrer T, Furnrohr BG, Niederweis M, Bilyy R, Schett G, Herrmann M, Berens C. The progression of cell death affects the rejection of allogeneic tumors in immune-competent mice—Implications for cancer therapy. *Front Immunol.* 2014;5:560.