

Overexpression of Long Noncoding RNA PTENP1 Inhibits Cell Proliferation and Migration via Suppression of miR-19b in Breast Cancer Cells

Xianbiao Shi, Xiaoqiao Tang, and Lei Su

Department of Breast Surgery, Nanjing Drum Tower Hospital, the Affiliated Hospital of Nanjing University Medical School, Nanjing, Jiangsu, P.R. China

This study aimed to investigate the effect of long noncoding RNA PTENP1 in the development of breast cancer (BC). Quantitative real-time PCR was utilized to determine the expression of PTENP1 in tissues and cell lines. pcDNA3.1 and shRNA were used to over- and low-express PTENP1 in BC cell lines, and miR-19b mimic and inhibitor were utilized to over- and low-express miR-19b. Then the abilities of cell survival, apoptosis, migration, and invasion were assessed in BC cells with different expression levels of PTENP1 and miR-19b. The expression of PTENP1 was significantly downregulated in both BC tissues and cell lines. Overexpressed PTENP1 could significantly increase cell survival, colony forming, migration, and invasion but decrease apoptosis in BC cell lines. However, overexpressed miR-19b performed contrary effects compared with PTENP1 on cell survival, colony forming, migration, invasion, and apoptosis in BC cell lines. miR-19b can be downregulated by PTENP1, and the effect of overexpressed PTENP1 on the PI3k/Akt pathway could be aborted by overexpressed miR-19b. PTENP1 performed a negative role in the development of BC via downregulating miR-19 probably through the PTEN/PI3K/Akt pathway.

Key words: Phosphatase and tensin homolog pseudogene 1 (PTENP1); Breast cancer (BC); MicroRNA-19b (miR-19b); Cell viability; Migration

INTRODUCTION

Breast cancer (BC) is the most common malignancy and accounts for the second highest number of cancer deaths worldwide in women¹. Recently, because of the popularization of the Westernized lifestyle in China, the diagnosis of BC increased significantly, especially in the obese and physically inactive population². It is estimated that the prevalence of BC was 268,600 cases, and mortality was expected to reach 69,500 cases in Chinese women for 2015³. Undoubtedly, these increased trends also induced a heavy burden on the economy and health of both patients and families⁴. Surgery resection, radiotherapy, and chemotherapy are the most common treatments for BC. However, because of their limitations, the clinical results and prognosis are still not satisfactory⁵. In recent decades, several biomarkers and targeted gene treatment genes have been introduced for BC therapy, but the clinical effect has still not improved owing to the comprehensive mechanism of BC development^{6,7}. Therefore, it is of great importance to further explore the mechanism of BC, so that we can obtain some new insights into the progression and treatment of BC.

Long noncoding RNAs (lncRNAs) are a class of transcripts more than 200 nucleotides in length, with limited protein-coding potential⁸. Recently, lncRNAs were identified to participate in several biological processes, such as the proliferation, migration, invasion, and apoptosis of cells^{9,10}. The investigation of lncRNAs in BC is currently ongoing¹¹. Singh et al. have documented that lncRNA-21A can serve as a potential tumor suppressor against the invasion of BC tumor cells¹². Long intergenic noncoding RNA for kinase activation (LINK-A) lncRNA was shown to activate a LINK-A-dependent pathway in correlation with hypoxia inhibitory factor 1 α (HIF1 α) to promote glycolysis reprogramming and tumorigenesis in BC by Lin et al.¹³. Growth arrest-specific 5 (GAS5) lncRNA can induce apoptosis in BC cells after responding to hormones¹⁴. Taken together, these findings suggest that lncRNAs play an important role in the genesis and progression of BC.

Lately, low levels of lncRNA phosphatase and tensin homolog pseudogene 1 (PTENP1) have been reported to significantly correlate with poor survival in patients with head and neck squamous cell carcinoma, and decreased

PTENP1 will promote the malignant behaviors of cancer cells¹⁵. Dong et al. have reported that PTENP1 can act as a biomarker to distinguish patients with gastric cancer from healthy individuals¹⁶. However, the biofunction of PTENP1 in BC has not been reported. Hence, a series of BC tissue samples and cell lines were collected in this study to primarily explore the mechanism of PTENP1 in BC. With these analyses, we aimed to obtain some basic information about PTENP1, so that we can provide some useful information for clinical diagnosis and treatment.

MATERIALS AND METHODS

Human Tissue Samples and Cell Lines

After surgical resection, a total of 20 tissue samples were collected from patients with BC at the Affiliated Drum Tower Hospital of Nanjing University Medical School (Jiangsu, P.R. China) from 2014 to 2016. All samples, including BC and corresponding adjacent normal tissues, were preserved in liquid nitrogen within 30 min after resection. All patients were informed of the procedures and signed consent forms. Normal human mammary epithelial cell line MCF10A and human BC cell lines MCF-7 and MDA-MB-231 (ATCC numbers: American Type HTB-22 and HTB-26, respectively; Manassas, VA, USA) were used in this study and cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco, Carlsbad, CA, USA) with 10% heat-inactivated fetal bovine serum (FBS; Gibco). The cells were maintained at 37°C in a humidified atmosphere of 5% CO₂ and passaged every 2–3 days to maintain exponential growth.

Quantitative Real-Time Reverse Transcription PCR (qRT-PCR)

Total RNA was extracted using TRIzol (Invitrogen, Carlsbad, CA, USA), and 1 mg of the total RNA was reverse transcribed to cDNA using the Omniscript RT Kit (Qiagen, Hilden, Germany). Each PCR mixture contained the following: 1 µl of Fast SYBR® Green Master Mix (Applied Biosystems, Foster City, CA, USA), 0.3 mM of each primer, and 50 ng of cDNA. Negative controls were performed using the same reaction mixtures without cDNA. Real-time PCR analyses were performed in StepOnePlus™ (Applied Biosystems) using the following conditions: 20 s at 95°C, and then 40 cycles of 3 s at 95°C and 30 s at 60°C. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as internal reference for the relative quantification analysis. Primers used in the measurement of PTENP1 were for the nonhomologous 3'-untranslated regions (3'-UTRs) as described previously¹⁷. The mature microRNAs (miRNAs) were quantified using the TaqMan MicroRNA Assays Kit (Applied

Biosystems) as described¹⁸, based on the quantitative stem loop PCR with U6 RNA as an internal control.

Western Blot

Total cellular extracts were obtained after the cells were washed twice with Hanks' balanced salt solution and lysed in RIPA buffer. Equal amounts of protein extracts (50 µg of protein per lane) were separated by 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene fluoride (PVDF) membranes (Millipore, Boston, MA, USA). The membranes were blocked with 5% skim milk in Tris-buffered saline with Tween (TBST) for 1 h. The membranes were incubated overnight at 4°C with primary antibodies to proteins (1:1,000 dilution) and GAPDH (1:1,000), and then a secondary antibody conjugated with horseradish peroxidase (1:5,000; ab7090) was added to culture the membrane continuously for 1 h at 37°C. The bands in the membranes were developed using enhanced chemiluminescence reagent (Santa Cruz) after washing for three times with 1× TBST. All antibodies were purchased from Abcam (Cambridge, UK), and their basic information is shown below: antibodies for B-cell CLL/lymphoma 2 (Bcl-2; ab119506), Bcl-2-associated X protein (Bax; ab77566), pro- and cleaved caspase 3 (ab32150 and ab2302), pro- and cleaved caspase 9 (ab135544 and ab2324), epithelial (E)-cadherin (ab1416), β-catenin (ab32572), vimentin (ab16051), α-smooth muscle actin (α-SMA; ab5831), phosphatase and tensin homolog (PTEN; ab228466), pyruvate dehydrogenase kinase 1 (PDK-1) (ab208187), phosphorylated phosphatidylinositol 3-kinase (p-PI3K), PI3K (ab125633), phosphorylated protein kinase B (p-Akt), and Akt (ab8804). GAPDH (ab181603) was used as a loading control.

Cell Transfection

Plasmid cDNA-PTENP1 was constructed by introducing a *Bam*HI–*Eco*RI fragment containing the PTENP1 cDNA into the same site in pcDNA3.1. The high-expressed and low-expressed PTENP1 were transfected with pcDNA-PTENP1 and short hairpin RNA for PTENP1 (sh-PTENP1), respectively. The pcDNA-PTENP1, pcDNA-control, sh-PTENP1 or sh-negative control (NC), miR-19b mimic, mimic NC, inhibitor NC, or miR-19b inhibitor was transfected into MCF-7 and/or MDA-MB-231 cells. Transfection was performed using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions.

Cell Proliferation Assays

Cell proliferation was determined using a Cell Proliferation Reagent Kit I (MTT) (Roche Applied Science, Penzberg, Germany). The MCF-7 and MDA-MB-231

cells transfected with pcDNA-PTENP1 were grown in 96-well plates. Cell proliferation was assessed every 24 h following the manufacturer's protocol. All experiments were performed in quadruplicate. For colony formation assays, a total of 500 cells for each cell line were placed in six-well plates and maintained in complete media. The medium was replaced every 4 days; 2 weeks later, cells were fixed with methanol and stained with 0.1% crystal violet (Sigma-Aldrich, St. Louis, MO, USA). Visible colonies were then counted. For each treatment group, wells were assessed in triplicate.

Cell Apoptosis

The MCF-7 and MDA-MB-231 cells transfected with pcDNA-PTENP1 were harvested 48 h after transfection by trypsinization. After being double stained with fluorescein isothiocyanate (FITC)-annexin V and propidium iodide (PI) using the FITC-Annexin-V Apoptosis Detection Kit (BD Biosciences, San Jose, CA, USA), the cells were analyzed with a flow cytometer (FACScan; BD Biosciences) equipped with CellQuest software (BD Biosciences). Cells were discriminated into viable cells, dead cells, early apoptotic cells, and apoptotic cells, and then the relative ratio of early apoptotic cells was compared with control from each experiment.

Cell Migration and Invasion Assays

For the migration assays, at 48 h posttransfection, 5×10^4 cells in serum-free medium were placed into the upper chamber of an insert (8-mm pore size; Millipore, Billerica, MA, USA). For the invasion assays, 1×10^5 cells in serum-free medium were placed into the upper chamber of an insert coated with Matrigel. Medium

containing 10% FBS was added to the lower chamber. After incubation for 24 h, the upper medium was discarded, and the upper membrane surface was wiped with cotton wool. Cells that had migrated or invaded through the membrane (lower membrane surface) were stained with methanol and 0.1% crystal violet, and then the staining results were imaged, and cell numbers were counted using an IX71 inverted microscope (Olympus, Tokyo, Japan). Experiments were independently repeated three times.

Luciferase Assay

The target relationship between lncRNA PTENP1 and miR-19b was evaluated using a dual-luciferase reporter assay. The full-length 3'-UTR of PTENP1 containing the miR-19b binding site was inserted into the *Xba*I site of the pGL3 vector (Promega, Madison, WI, USA) to construct the pGL3-PTENP1-3'-UTR-WT reporter. The pGL3-PTENP1-3'-UTR-MUT reporter, containing a mutated PTENP1 binding sequence, was synthesized using a site-directed mutagenesis kit (Stratagene, La Jolla, CA, USA). Afterward, cells (1×10^6) were transfected with 1 μ g of the silenced PTENP1 vector (or control) and a *Renilla* luciferase expression vector using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. After 36 h of transfection, luciferase activity was measured using the Dual-Luciferase[®] Reporter Assay System (Promega). *Renilla* luciferase activity served as an internal control.

Statistical Analysis

Differences between groups were analyzed using Student's *t*-test or chi-square test. Statistical analyses were




Figure 1. Long noncoding RNA (lncRNA) phosphatase and tensin homolog pseudogene 1 (PTENP1) was decreased in breast cancer (BC) tissues and cells as determined by quantitative real-time PCR. (A) Expression of PTENP1 in tissue samples. ***p* < 0.01 compared to nontumor. (B) Expression of PTENP1 in MCF10A, MCF-7, and MDA-MB-231. ***p* < 0.01 compared to MCF10A.




Figure 2. Overexpressed PTENP1 inhibited cell proliferation and promoted cell apoptosis. (A) Relative expression of PTENP1 determined by quantitative real-time PCR. (B) Effect of PTENP1 overexpression on cell viability in MCF-7 cells. (C) Effect of PTENP1 overexpression on cell viability in MDA-MB-231 cells. (D) Effect of PTENP1 overexpression on colony formation in both MCF-7 and MDA-MB-231 cells. (E) Influence of PTENP1 overexpression on apoptosis in both MCF-7 and MDA-MB-231. (F) Expression of apoptosis markers determined by Western blotting in both MCF-7 and MDA-MB-231. Bcl-2, B-cell lymphoma 2; Bax, Bcl-2-associated X protein; GAPDH, glyceraldehyde 3-phosphate dehydrogenase. * $p < 0.05$ and ** $p < 0.01$ compared to pcDNA-control.

performed using SPSS version 18.0 (SPSS, Chicago, IL, USA). For all statistical analyses, a value of $p < 0.05$ was considered statistically significant.

RESULTS

Expressions of PTENP1 in BC Tissues and Cell Lines

In the current study, a total of 20 patients were enrolled, and the expressions of PTENP1 in both tumor and nontumor tissues were examined via qRT-PCR. According to the results of qRT-PCR, a significantly lower expression of PTENP1 was identified in tumor tissue compared with nontumor tissue (Fig. 1A). Meanwhile, expression levels of PTENP1 in BC cell lines MCF-7 and MDA-MB-231 were examined by qRT-PCR with the MCF10A as the normal control. Similar to the result in tissue, a significant reduction was obtained in MCF-7 and MDA-MB-231 compared with MCF10A cells (Fig. 1B).

Overexpression of PTENP1 on Cell Proliferation and Cell Apoptosis

For further analysis, PTENP1 was overexpressed in MCF-7 and MDA-MB-231 cells (Fig. 2A). Then the influence of overexpressed PTENP1 on cell proliferation and apoptosis was assessed. The MTT result showed that overexpressed PTENP1 significantly inhibited the proliferation of MCF-7 and MDA-MB-231 at 48 and 72 h (Fig. 2B and C). Overexpressed PTENP1 also could significantly suppress the colony-forming abilities of MCF-7 and MDA-MB-231 (Fig. 2D). The degree of apoptosis in MCF-7 and MDA-MB-231 cell lines was also investigated. The results showed that overexpressed PTENP1 significantly increased apoptosis in both MCF-7

and MDA-MB-231 cells (Fig. 2E). Meanwhile, examination of apoptosis factors also revealed that overexpressed PTENP1 significantly increased the expression levels of Bax, cleaved caspase 3, and cleaved caspase 9, but remarkably decreased the expressions of Bcl-2, procaspase 3, and procaspase 9 (Fig. 2F).

Overexpression of PTENP1 on Cell Migration and Invasion

The influence of PTENP1 on cell migration and invasion was also investigated. The results revealed that overexpressed PTENP1 could significantly inhibit the migrations of MCF-7 and MDA-MB-231 cells (Fig. 3A). Meanwhile, overexpressed PTENP1 also can suppress the invasion abilities of MCF-7 and MDA-MB-231 cells (Fig. 3B).

miR-19b Is a Target Gene of PTENP1

To further reveal the possible mechanism of action, the downstream target of PTENP1 was explored (Fig. 4A). Based on the luciferase assay, miR-19b was identified to be negatively regulated by PTENP1 in MCF-7 cells (Fig. 4B and C). The expression of miR-19b in BC tissue and cell lines was also examined. The results showed that the expression of miR-19b was significantly increased in BC tissue compared with nontumor tissue (Fig. 4D). Significantly increased expression levels were also detected in the MCF-7 and MDA-MB-231 cell lines compared with MCF10A (Fig. 4E).

miR-19b Promotes Cell Proliferation and Metastasis

Moreover, the biofunction of miR-19b in BC was explored. The effect on miR-19b levels of overexpression and knockdown of miR-19b vectors in MCF-7 was




Figure 3. PTENP1 overexpression inhibited cell migration and invasion. (A) Effect of PTENP1 overexpression on migration assay. (B) Effect of PTENP1 overexpression on invasion assay. ** $p < 0.01$ compared to pcDNA-control.




Figure 4. MicroRNA-19b (miR-19b) was a target gene of PTENP1. (A) Schematic of the miR-19b putative binding site in wild-type PTENP1. (B) Luciferase activity assay performed between PTENP1 and miR-19b. WT, wild type; MUT, mutant. * $p < 0.05$ compared to pre-negative control (NC). (C) Expression of miR-19b in MCF-7 with different expression of PTENP1. sh, short hairpin. * $p < 0.05$ compared to sh. (D) Expression of miR-19b in tissue. *** $p < 0.01$ compared to nontumor. (E) Relative expression of miR-19b in cancer cell lines. ** $p < 0.001$ compared to MCF10A.

assessed in Figure 5A. The results demonstrated that overexpression of miR-19b can significantly increase the ability of survival, but inhibition of miR-19b would decrease the ability of survival (Fig. 5B). The same effects were also observed on colony forming (Fig. 5C), migration (Fig. 5D), and invasion (Fig. 5E) in MCF-7 cells. Results of Western blotting also showed that overexpressed miR-19b could significantly downregulate the expression levels of E-cadherin and β -catenin but upregulate the expression levels of vimentin and α -SMA, while inhibiting the expression of miR-19b would markedly upregulate the expression levels of E-cadherin

and β -catenin but downregulate the expression levels of vimentin and α -SMA (Fig. 5F). In addition, upregulating the expression of miR-19b decreased the apoptosis of MCF-7, but downregulating the expression of miR-19b remarkably increased the apoptosis of MCF-7 cells (Fig. 5G). Moreover, the Western blot also showed that increasing the expression of miR-19b would significantly increase the expression of procaspase 3 and procaspase 9 but decrease the expression of Bax-2, cleaved caspase 3, and cleaved caspase 9, whereas suppressing the expression of miR-19b would markedly inhibit the expression of Bcl-2, procaspase 3, and procaspase 9, but promote




Figure 5. miR-19b promoted cell proliferation and metastasis and inhibited cell apoptosis. (A) Relative expression of miR-19b following transfection with mimics or inhibitors, as determined by quantitative real-time PCR. (B) Effect of miR-19b mimics and inhibitors on cell viability. (C) Effect of miR-19b mimics and inhibitors on colony forming. (D) Effect of miR-19b mimics and inhibitors on migration assay. (E) Effect of miR-19b mimics and inhibitors on invasion assay. (F) Effect of miR-19b mimics and inhibitors on expression of migration markers as determined by Western blotting. (G) Effect of miR-19b mimics and inhibitors on apoptosis. (H) Effect of miR-19b mimics and inhibitors on expression of apoptosis markers as determined by Western blotting. E-cadherin, epithelial-cadherin; α-SMA, α-smooth muscle actin. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared to equivalent NC.

the expression of Bax-2, cleaved caspase 3, and cleaved caspase 9 (Fig. 5H).

PTENP1 Inhibits BC Cell Proliferation and Metastasis by Downregulation of miR-19b

The correlation between PTENP1 and miR-19b was also further investigated. The analytical result showed

that overexpression of PTENP1 presented obvious inhibitory effects on MCF-7 cell viability (Fig. 6A), colony forming (Fig. 6B), migration (Fig. 6C), and invasion (Fig. 6D) and expression of migration proteins (Fig. 6E). However, these results could be significantly opposed by overexpression of miR-19b. Moreover, overexpression of PTENP1 also could conduct a significant promotive




Figure 6. PTENP1 inhibited BC cell proliferation and metastasis by downregulation of miR-19b. (A) Effect of overexpression of PTENP1 and miR-19b on cell viability. (B) Effect of miR-19b mimics and inhibitors on colony formation. (C) Effect of miR-19b mimics and inhibitors on migration. (D) Effect of miR-19b mimics and inhibitors on invasion. (E) Effect of miR-19b mimics and inhibitors on expression of migration markers determined by Western blotting. (F) Effect of miR-19b mimics and inhibitors on apoptosis. (G) Effect of miR-19b mimics and inhibitors on expression of apoptosis markers determined by Western blotting. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared with the nontumor group.

effect on apoptosis of MCF-7 (Fig. 6F), and the apoptotic factors were also changed correspondingly (Fig. 6G).

Effects of PTENP1 on the PTEN/PI3K/Akt Pathway

Finally, the possible PTENP1 pathway that might be involved in BC genesis and progression was further explored. The results showed that overexpression of PTENP1 could significantly increase the expression of PTEN, while it could decrease the expression of PDK-1, p-PI3K, PI3K, and p-Akt. However, overexpression of miR-19b could decrease the expression of PTEN and PI3K and increase the expression of PDK-1, p-PI3K, and p-Akt. Moreover, the influence of PTENP1 on the PI3K/Akt pathway can be significantly aborted by the overexpression of miR-19b (Fig. 7).

DISCUSSION

In the current study, PTENP1 was significantly downregulated in both BC tissue samples and cell lines. Meanwhile, upregulation of PTENP1 significantly inhibited cell viability, colony forming, migration, and invasion and promoted cell apoptosis in BC cell lines. miR-19b is an important downstream target of PTENP1 and may serve a positive role in the development of BC. Further

analyses suggested that the effect of the low expression of PTENP1 could be reversed by inhibiting the expression of miR-19b through the PTEN/PI3K/Akt pathway.

PTENP1 originates from a duplication of the tumor suppressor PTEN¹⁷. As a pseudogene of PTEN, PTENP1 has several binding sites for miRNAs that target PTEN in the 3'-UTR¹⁸. Owing to this manifestation, PTENP1 also presents an important role against the development of tumors. Tay et al. have documented that overexpressed PTENP1 can significantly inhibit cell growth via competing with PTEN¹⁹. A previous study had reported that the expression levels of PTEN and PTENP1 were both downregulated in hepatocellular carcinoma (HCC), and overexpressed PTENP1 will significantly impair the tumorigenic ability of HCC cell lines and increase the autophagy and apoptosis of cell lines²⁰. Downregulated PTENP1 expression was also observed in BC tissue samples and cell lines. Overexpressed PTENP1 negatively regulated cell viability, proliferation, migration, and invasion of BC cell lines. This evidence suggested that PTENP1 acted via a negative role in the development of BC. However, the mechanism for the downregulation of PTENP1 is not clearly elucidated. Kovalenko et al. have indicated that the methylation of PTENP1 in its 5'-terminal regions contributes to this decrease in expression²¹. Thus, we deduced that the methylation of PTENP1 might be the explanation for the decline of PTENP1 expression, but this still needs to be verified.

PTEN can be regulated by several mechanisms, including miRNAs²². miR-19b has been confirmed to target PTEN through binding sites in the 3'-UTR, which is also contained in PTENP1²³. Therefore, the biofunction of miR-19b was also explored in this study. With an increased expression of miR-19b, significant promotive effects were identified on cell viability, colony forming, migration, and invasion, and an obvious inhibitive effect on apoptosis in BC cell lines, while these effects can be opposed by downregulating the expression of miR-19b. These results meant that miR-19b served a positive role in the progression of BC. This positive role has also been identified in colon cancer²⁴ and non-small cell lung cancer²⁵. Downregulating miR-19b expression with an inhibitor appeared to reverse the effect of low PTENP1 expression on cell viability, colony forming, migration, and invasion. All of these results indicate that miR-19b is a downstream target of PTENP1. The PI3K/Akt pathway is a crucial role for PTEN to develop its biofunction. Chen et al. have revealed that elevated expression levels of PTENP1 and PTEN can suppress cell proliferation, migration, and invasion via inhibiting the PI3K/Akt pathway in HCC²⁰. Hence, the PI3K/Akt pathway was also explored in this study, and the results showed that increasing PTENP1 expression could also increase




Figure 7. Expression of proteins involved in the PI3K/Akt pathway was changed by aberrant expression of PTENP1 and miR-19b. PTEN, phosphatase and tensin homolog; PDK-1, pyruvate dehydrogenase kinase 1; p-PI3K, phosphorylated phosphatidylinositol 3-kinase; p-Akt, phosphorylated protein kinase B.

the expression of PTEN but reduce the activation of the PDK-1 and PI3K/Akt pathway, while miR-19b acted in a contrary way. Taken together, the PTEN/PI3K/Akt pathway was the main signal transduction pathway for PTENP1 in BC.

In conclusion, PTENP1 is an important tumor suppressor for the development of BC, but miR-19b is an oncogenic gene for BC. PTENP1 can inhibit cell viability, colony forming, migration, and invasion and promote cell apoptosis via downregulating miR-19 potentially through the PTEN/PI3K/Akt pathway. Therefore, PTENP1 can serve as a potential target gene for BC treatment.

ACKNOWLEDGMENTS: *The authors declare no conflicts of interest.*

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin.* 2016;66(1):7–30.
2. Varghese C, Shin HR. Strengthening cancer control in China. *Lancet Oncol.* 2014;15(5):484–5.
3. Chen W, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, Yu XQ, He J. Cancer statistics in China, 2015. *CA Cancer J Clin.* 2016;66(2):115–32.
4. Chen W. Cancer statistics: Updated cancer burden in China. *Chinese J Cancer Res.* 2015;27(1):1.
5. McGale P, Taylor C, Correa C, Cutter D, Duane F, Ewertz M, Gray R, Mannu G, Peto R, Whelan T. Effect of radiotherapy after mastectomy and axillary surgery on 10-year recurrence and 20-year breast cancer mortality: Meta-analysis of individual patient data for 8135 women in 22 randomised trials. *Lancet* 2014;383(9935):2127–35.
6. Söletormos G, Duffy MJ, Othman AHS, Verheijen RH, Tholander B, Jr BR, Gaarenstroom KN, Sturgeon CM, Bonfrer JM, Petersen PH. Clinical use of cancer biomarkers in epithelial ovarian cancer: Updated guidelines from the European Group on Tumor Markers. *Int J Gynecol Cancer* 2017;75(1):284–98.
7. Early Breast Cancer Trialists' Collaborative Group. Aromatase inhibitors versus tamoxifen in early breast cancer: Patient-level meta-analysis of the randomised trials. *Lancet* 2015;386(10001):1341–52.
8. Shi SJ, Wang LJ, Yu B, Li YH, Jin Y, Bai XZ. LncRNA-ATB promotes trastuzumab resistance and invasion-metastasis cascade in breast cancer. *Oncotarget* 2015;6(13):11652–63.
9. Sun YW, Chen YF, Li J, Huo YM, Liu DJ, Hua R, Zhang JF, Liu W, Yang JY, Fu XL. A novel long non-coding RNA ENST00000480739 suppresses tumour cell invasion by regulating OS-9 and HIF-1 α in pancreatic ductal adenocarcinoma. *Br J Cancer* 2014;111(11):2131–41.
10. Meng J, Li P, Zhang Q, Yang Z, Fu S. A four-long non-coding RNA signature in predicting breast cancer survival. *J Exp Clin Cancer Res.* 2014;33(1):84.
11. Mendell JT. Targeting a long noncoding RNA in breast cancer. *N Engl J Med.* 2016;374(23):2287–9.
12. Singh R, Mo Y. LncRNA-21a is a potential tumor suppressor in breast cancer. *Cancer Res.* 2014;74(19 Suppl):3555.
13. Lin A, Li C, Xing Z, Hu Q, Liang K, Han L, Wang C, Hawke DH, Wang S, Zhang Y. The LINK-A lncRNA activates normoxic HIF1 [alpha] signalling in triple-negative breast cancer. *Nat Cell Biol.* 2016;18(2):213–24.
14. Pickard MR, Williams GT. The hormone response element mimic sequence of GAS5 lncRNA is sufficient to induce apoptosis in breast cancer cells. *Oncotarget* 2016;7(9):10104–16.
15. Liu J, Xing Y, Xu L, Chen W, Cao W, Zhang C. Decreased expression of pseudogene PTENP1 promotes malignant behaviours and is associated with the poor survival of patients with HNSCC. *Sci Rep.* 2017;7:41179.
16. Dong L, Qi P, Xu MD, Ni SJ, Huang D, Xu QH, Weng WW, Tan C, Sheng WQ, Zhou XY. Circulating CUDR, LSINCT-5 and PTENP1 long non-coding RNAs in sera distinguish patients with gastric cancer from healthy controls. *Int J Cancer* 2015;133(5):1128–35.
17. Xu J, Zhang J. Are human translated pseudogenes functional? *Mol Biol Evol.* 2016;33(3):755–60.
18. Swami M. Small RNAs: Pseudogenes act as microRNA decoys. *Nat Rev Cancer* 2010;10(8):530–1.
19. Tay Y, Kato J, Salmena L, Weiss D, Tan SM, Ala U, Karreth F, Poliseno L, Provero P, Cunto FD. Coding-independent regulation of the tumor suppressor PTEN by competing endogenous mRNAs. *Cell* 2011;147(2):344–57.
20. Chen CL, Tseng YW, Wu JC, Chen GY, Lin KC, Hwang SM, Hu YC. Suppression of hepatocellular carcinoma by baculovirus-mediated expression of long non-coding RNA PTENP1 and microRNA regulation. *Biomaterials* 2015;44:71–81.
21. Kovalenko TF, Sorokina AV, Ozolinya LA, Patrushev LI. Methylation of the pseudogene PTENP1 5'-terminal region in endometrial cancer and hyperplasia. *Russ J Bioorg Chem.* 2013;39(4):397–405.
22. He L. Post-transcriptional regulation of PTEN dosage by non-coding RNAs. *Sci Signal* 2010;3(146):pe39.
23. Olive V, Bennett MJ, Walker JC, Ma C, Jiang I, Cordon-Cardo C, Li Q-J, Lowe SW, Hannon GJ, He L. miR-19 is a key oncogenic component of mir-17-92. *Gene Dev.* 2009;23(24):2839–49.
24. Kurokawa K, Tanahashi T, Iima T, Yamamoto Y, Akaike Y, Nishida K, Masuda K, Kuwano Y, Murakami Y, Fukushima M. Role of miR-19b and its target mRNAs in 5-fluorouracil resistance in colon cancer cells. *J Gastroenterol.* 2012;47(8):883–95.
25. Wu C, Cao Y, He Z, He J, Hu C, Duan H, Jiang J. Serum levels of miR-19b and miR-146a as prognostic biomarkers for non-small cell lung cancer. *Tohoku J Exp Med.* 2014;232(2):85–95.